




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/63, 9/88, C07R 16/40		A2	(11) International Publication Number: WO 00/24913
			(43) International Publication Date: 4 May 2000 (04.05.00)
(21) International Application Number: PCT/US99/24879 (22) International Filing Date: 22 October 1999 (22.10.99) (30) Priority Data: 09/177,776 23 October 1998 (23.10.98) US 09/178,115 23 October 1998 (23.10.98) US (71) Applicant (for AM BF BJ CF CG CI CM GA GN GW ML MR NE SN TD TG only): BAYER CORPORATION [US/US]; 63 North Street, Medfield, MA 02052 (US). (71) Applicant (for all designated States except US): INSTITUTE OF VIROLOGY [SK/SK]; Slovak Academy of Sciences, Dubravská Cesta 9, 842 46 Bratislava (SK). (72) Inventors; and (75) Inventors/Applicants (for US only): ZAVADA, Jan [CZ/CZ]; Na pekne vyhlidce 1, 160 00 Prague 6 (CZ). PASTOREKOVA, Silvia [SK/SK]; I. Bukovciana 18, 841 07 Bratislava (SK). PASTOREK, Jaromir [SK/SK]; I. Bukovciana 18, 841 07 Bratislava (SK). (74) Agent: LAUDER, Leona, L.; Suite 610, 369 Pine street, San Francisco, CA 94104-3313 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.	

(54) Title: MN GENE AND PROTEIN



(57) Abstract

Identified herein is the location of the MN protein binding site, and MN proteins/polypeptides that compete for attachment to vertebrate cells with immobilized MN protein. Such MN proteins/polypeptides prevent cell-cell adhesion and the formation of intercellular contacts. The MN protein binding site is a therapeutic target that can be blocked by organic or inorganic molecules, preferably organic molecules, more preferably proteins/polypeptides that specifically bind to that site. Therapeutic methods for inhibiting the growth of preneoplastic/neoplastic vertebrate cells that abnormally express MN protein are disclosed. Vectors are provided that encode the variable domains of MN-specific antibodies and a flexible linker polypeptide separating those domains. Further vectors are disclosed that encode a cytotoxic protein/polypeptide operatively linked to the MN gene promoter, and which vectors preferably further encode a cytokine. The MN gene promoter is characterized, and the binding site for a repressor of MN transcription is disclosed.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

MN GENE AND PROTEIN
FIELD OF THE INVENTION

The present invention is in the general area of medical genetics and in the fields of biochemical engineering, immunochemistry and oncology. More specifically, it relates to the MN gene – a cellular gene considered to be an oncogene, which encodes the oncoprotein now known alternatively as the MN protein, the MN/CA IX isoenzyme or the MN/G250 protein.

BACKGROUND OF THE INVENTION

Zavada et al., International Publication Number WO 93/18152 (published 16 September 1993) and U.S. Patent No. 5,387,676 (issued February 7, 1996), describe the elucidation of the biological and molecular nature of MaTu which resulted in the discovery of the MN gene and protein. The MN gene was found to be present in the chromosomal DNA of all vertebrates tested, and its expression to be strongly correlated with tumorigenicity.

The MN protein was first identified in HeLa cells, derived from a human carcinoma of cervix uteri. It is found in many types of human carcinomas (notably uterine cervical, ovarian, endometrial, renal, bladder, breast, colorectal, lung, esophageal, and prostate, among others). Very few normal tissues have been found to express MN protein to any significant degree. Those MN-expressing normal tissues include the human gastric mucosa and gallbladder epithelium, and some other normal tissues of the alimentary tract. Paradoxically, MN gene expression has been found to be lost or reduced in carcinomas and other preneoplastic/neoplastic diseases in some tissues that normally express MN, e.g., gastric mucosa.

In general, oncogenesis may be signified by the abnormal expression of MN protein. For example, oncogenesis may be signified: (1) when MN protein is present in a tissue which normally does not express MN protein to any significant degree; (2) when MN protein is absent from a tissue that normally expresses it; (3) when MN gene expression is at a significantly increased level, or at a significantly reduced

level from that normally expressed in a tissue; or (4) when MN protein is expressed in an abnormal location within a cell.

Zavada et al., WO 93/18152 and Zavada et al., WO 95/34650 (published 21 December 1995) disclose how the discovery of the MN gene and protein and the strong association of MN gene expression and tumorigenicity led to the creation of methods that are both diagnostic/prognostic and therapeutic for cancer and precancerous conditions. Methods and compositions were provided therein for identifying the onset and presence of neoplastic disease by detecting or detecting and quantitating abnormal MN gene expression in vertebrates. Abnormal MN gene expression can be detected or detected and quantitated by a variety of conventional assays in vertebrate samples, for example, by immunoassays using MN-specific antibodies to detect or detect and quantitate MN antigen, by hybridization assays or by PCR assays, such as RT-PCR, using MN nucleic acids, such as, MN cDNA, to detect or detect and quantitate MN nucleic acids, such as, MN mRNA.

Zavada et al, WO 93/18152 and WO 95/34650 describe the production of MN-specific antibodies. A representative and preferred MN-specific antibody, the monoclonal antibody M75 (Mab M75), was deposited at the American Type Culture Collection (ATCC) in Manassus, VA (USA) under ATCC Number HB 11128. The M75 antibody was used to discover and identify the MN protein and can be used to identify readily MN antigen in Western blots, in radioimmunoassays and immunohistochemically, for example, in tissue samples that are fresh, frozen, or formalin-, alcohol-, acetone- or otherwise fixed and/or paraffin-embedded and deparaffinized. Another representative and preferred MN-specific antibody, Mab MN12, is secreted by the hybridoma MN 12.2.2, which was deposited at the ATCC under the designation HB 11647. Example 1 of Zavada et al., WO 95/34650 provides representative results from immunohistochemical staining of tissues using MAb M75, which results support the designation of the MN gene as an oncogene.

Many studies have confirmed the diagnostic/prognostic utility of MN. The following articles discuss the use of the MN-specific MAb M75 in diagnosing/prognosing precancerous and cancerous cervical lesions: Leff, D. N., "Half a Century of HeLa Cells: Transatlantic Antigen Enhances Reliability of Cervical Cancer Pap Test, Clinical Trials Pending," BioWorld® Today: The Daily Biotechnology

Newspaper, 9(55) (March 24, 1998); Stanbridge, E. J., "Cervical marker can help resolve ambiguous Pap smears," Diagnostics Intelligence, 10(5): 11 (1998); Liao and Stanbridge, "Expression of the MN Antigen in Cervical Papanicolaou Smears Is an Early Diagnostic Biomarker of Cervical Dysplasia," Cancer Epidemiology, Biomarkers & Prevention, 5: 549-557 (1996); Brewer et al., "A Study of Biomarkers in Cervical Carcinoma and Clinical Correlation of the Novel Biomarker MN," Gynecologic Oncology, 63: 337-344 (1996); and Liao et al., "Identification of the MN Antigen as a Diagnostic Biomarker of Cervical Intraepithelial Squamous and Glandular Neoplasia and Cervical Carcinomas," American Journal of Pathology, 145(3): 598-609 (1994).

10 Premalignant and Malignant Colorectal Lesions. MN has been detected in normal gastric, intestinal, and biliary mucosa. [Pastorekova et al., Gastroenterology, 112: 398-408 (1997).] Immunohistochemical analysis of the normal large intestine revealed moderate staining in the proximal colon, with the reaction becoming weaker distally. The staining was confined to the basolateral surfaces of the cryptal epithelial
15 cells, the area of greatest proliferative capacity. As MN is much more abundant in the proliferating cryptal epithelium than in the upper part of the mucosa, it may play a role in control of the proliferation and differentiation of intestinal epithelial cells. Cell proliferation increases abnormally in premalignant and malignant lesions of the colorectal epithelium, and therefore, is considered an indicator of colorectal tumor
20 progression. [Risio, M., J. Cell Biochem, 16G: 79-87 (1992); and Moss et al., Gastroenterology, 111: 1425-1432 (1996).]

The MN protein is now considered to be the first tumor-associated carbonic anhydrase (CA) isoenzyme that has been described. Carbonic anhydrases (CAs) form a large family of genes encoding zinc metalloenzymes of great physiological
25 importance. As catalysts of reversible hydration of carbon dioxide, these enzymes participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, formation of aqueous humor, cerebrospinal fluid, saliva and gastric acid [reviewed in Dodgson et al., The Carbonic Anhydrases, Plenum Press, New York-London, pp. 398 (1991)]. CAs are widely distributed in different living
30 organisms.

In mammals, at least seven isoenzymes (CA I-VII) and a few CA-related proteins (CARP/CA VIII, RPTP- β , RPTP- τ) had been identified [Hewett-Emmett and

Tashian, Mol. Phyl. Evol., 5: 50-77 (1996)], when analysis of the MN deduced amino acid sequence revealed a striking homology between the central part of the MN protein and carbonic anhydrases, with the conserved zinc-binding site as well as the enzyme's active center. Then MN protein was found to bind zinc and to have CA activity. Based on that data, the MN protein is now considered to be the ninth carbonic anhydrase isoenzyme – MN/CA IX. [Opavsky et al., Genomics, 33: 480-487 (May 1996)]. [See also, Hewett-Emmett, supra, wherein CA IX is suggested as a nomenclatural designation.]

CAs and CA-related proteins show extensive diversity in both their tissue distribution and in their putative or established biological functions [Tashian, R. E., Adv. in Genetics, 30: 321-356 (1992)]. Some of the CAs are expressed in almost all tissues (CA II), while the expression of others appears to be more restricted (CA VI and CA VII in salivary glands). In cells, they may reside in the cytoplasm (CA I, CA II, CA III, and CA VII), in mitochondria (CA V), in secretory granules (CA VI), or they may associate with membrane (CA IV). Occasionally, nuclear localization of some isoenzymes has been noted [Parkkila et al., Gut, 35: 646-650 (1994); Parkkila et al., Histochem. J., 27: 133-138 (1995); Mori et al., Gastroenterol., 105: 820-826 (1993)].

The CAs and CA-related proteins also differ in kinetic properties and susceptibility to inhibitors [Sly and Hu, Annu. Rev. Biochem., 64: 375-401 (1995)]. In the alimentary tract, carbonic anhydrase activity is involved in many important functions, such as saliva secretion, production of gastric acid, pancreatic juice and bile, intestinal water and ion transport, fatty acid uptake and biogenesis in the liver. At least seven CA isoenzymes have been demonstrated in different regions of the alimentary tract. However, biochemical, histochemical and immunocytochemical studies have revealed a considerable heterogeneity in their levels and distribution [Swensen, E. R., "Distribution and functions of carbonic anhydrase in the gastrointestinal tract," In: The Carbonic Anhydrases. Cellular Physiology and Molecular Genetics, (Dodgson et al. eds.) Plenum Press, New York, pages 265-287 (1991); and Parkkila and Parkkila, Scan J. Gastroenterol., 31: 305-317 (1996)]. While CA II is found along the entire alimentary canal, CA IV is linked to the lower gastrointestinal tract, CA I, III and V are present in only a few tissues, and the expression of CA VI and VII is restricted to

salivary glands [Parkkila et al., Gut, 35: 646-650 (1994); Fleming et al., J. Clin. Invest., 96: 2907-2913 (1995); Parkkila et al., Hepatology, 24: 104 (1996)].

MN/CA IX has a number of properties that distinguish it from other known CA isoenzymes and evince its relevance to oncogenesis. Those properties include its density dependent expression in cell culture (e.g., HeLa cells), its correlation with the tumorigenic phenotype of somatic cell hybrids between HeLa and normal human fibroblasts, its close association with several human carcinomas and its absence from corresponding normal tissues [e.g., Zavada et al., Int. J. Cancer, 54: 268-274 (1993); Pastorekova et al., Virology, 187: 620-626 (1992); Liao et al., Am. J. Pathol., 145: 598-609 (1994); Pastorek et al., Oncogene, 9: 2788-2888 (1994); Cote, Women's Health Weekly: News Section, p. 7 (March 30, 1998); Liao et al., Cancer Res., 57: 2827 (1997); Vermylen et al., "Expression of the MN antigen as a biomarker of lung carcinoma and associated precancerous conditions," Proceedings AACR, 39: 334 (1998); McKiernan et al., Cancer Res., 57: 2362 (1997); and Turner et al., Hum. Pathol., 28(6): 740 (1997)]. In addition, the *in vitro* transformation potential of MN/CA IX cDNA has been demonstrated in NIH 3T3 fibroblasts [Pastorek et al., id.].

The MN protein has also been identified with the G250 antigen. Uemura et al., "Expression of Tumor-Associated Antigen MN/G250 in Urologic Carcinoma: Potential Therapeutic Target," J. Urol., 154 (4 Suppl.): 377 (Abstract 1475; 1997) states: "Sequence analysis and database searching revealed that G250 antigen is identical to MN, a human tumor-associated antigen identified in cervical carcinoma (Pastorek et al., 1994)."

SUMMARY OF THE INVENTION

25

Identified herein is the location of the MN protein binding site. Of particular importance is the region within the proteoglycan-like domain, aa 61-96 (SEQ ID NO: 97) which contains a 6-fold tandem repeat of 6 amino acids, and within which the epitope for the M75 MAb resides in at least two copies, and within which the MN binding site is considered to be located. An alternative MN binding site may be located in the CA domain.

30

Also identified are MN proteins and MN polypeptides that compete for attachment to cells with immobilized MN protein. Such MN proteins/polypeptides prevent cell-cell adhesion and the formation of intercellular contacts.

Disclosed herein are cell adhesion assay methods that are used to identify
5 binding site(s) on the MN protein to which vertebrate cells, preferably mammalian cells, more preferably human cells, bind. Such a MN binding site is then identified as a therapeutic target which can be blocked with MN-specific antibodies, or inorganic or organic molecules, preferably organic molecules, more preferably proteins/polypeptides that specifically bind to said site.

10 Further disclosed are therapeutic methods to treat patients with preneoplastic/neoplastic disease associated with or characterized by abnormal MN expression, which methods are based on blocking said MN binding site with molecules, inorganic or organic, but preferably organic molecules, more preferably proteins/polypeptides, that bind specifically to said binding site. The growth of a
15 vertebrate preneoplastic/neoplastic cell that abnormally expresses MN protein can be inhibited by administering such organic or inorganic molecules, preferably organic molecules, more preferably proteins/polypeptides in a therapeutically effective amount in a physiologically acceptable formulation. Such a preferred therapeutic protein/polypeptide is herein considered to comprise an amino acid sequence selected
20 from the group consisting of SEQ ID NOS: 107-109. Such heptapeptides are considered to be comprised by MN protein partner(s). Blocking the interaction between MN protein and its binding partner(s), is expected to lead to a decrease of tumor growth.

Further provided are other therapeutic methods wherein the growth of a
25 vertebrate, preferably mammalian, more preferably human, preneoplastic or neoplastic cell that abnormally expresses MN protein is inhibited. Said methods comprise transfecting said cell with a vector comprising an expression control sequence operatively linked to a nucleic acid encoding the variable domains of an MN-specific antibody, wherein said domains are separated by a flexible linker peptide, preferably
30 SEQ ID NO: 116. Preferably said expression control sequence comprises the MN gene promoter.

Still further therapeutic methods comprise transfecting said cell with a vector comprising a nucleic acid that encodes a cytotoxic protein/polypeptide, such as HSVtk, operatively linked to the MN gene promoter. Such a therapeutic vector may also comprise a nucleic acid encoding a cytokine, such as, IL-2 or IFN.

5 Aspects of the instant invention disclosed herein are described in more detail as follows. The therapeutic use of organic or inorganic molecules, preferably organic molecules, is disclosed. Preferred such molecules bind specifically to a site on MN protein to which vertebrate cells adhere in a cell adhesion assay, wherein said molecule when tested in vitro inhibits the adhesion of cells to MN protein. Further
10 preferred are such molecules, which when in contact with a vertebrate preneoplastic or neoplastic cell that abnormally expresses MN protein, inhibit the growth of said cell. Said vertebrate cells are preferably mammalian and more preferably human.

Preferably such a molecule is organic, and more preferably such a organic molecule is a protein or a polypeptide. Still further preferably, said protein or
15 polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 107, 108, 109, 137 and 138. Even more preferably, said polypeptide is selected from the group consisting of SEQ ID NOS: 107, 108, 109, 137 and 138.

The site on MN proteins to which vertebrate cells adhere in said cell adhesion assay is preferably within the proteoglycan-like domain [SEQ ID NO: 50] or
20 within the carbonic anhydrase domain [SEQ ID NO: 51] of the MN protein. Preferably that site comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 10 and 97-106. Still further preferably, that site has an amino acid sequence selected from the group consisting of SEQ ID NOS: 10 and 97-106.

Another aspect of this invention concerns MN proteins and MN
25 polypeptides which mediate attachment of vertebrate cells in a cell adhesion assay, wherein said MN protein or MN polypeptide when introduced into the extracellular fluid environment of vertebrate cells prevents the formation of intercellular contacts and the adhesion of said vertebrate cells to each other. Such MN proteins and MN polypeptides may be useful to inhibit the growth of vertebrate preneoplastic or
30 neoplastic cells that abnormally express MN protein, when such MN proteins or MN polypeptides are introduced into the extracellular fluid environment of such vertebrate cells. Said vertebrate cells are preferably mammalian, and more preferably human.

Said MN proteins or MN polypeptides which mediate attachment of vertebrate cells in a cell adhesion assay, preferably have amino acid sequences from SEQ ID NO: 97, from SEQ ID NO: 50, or from SEQ ID NO: 51, more preferably from SEQ ID NO: 50. Still more preferably such MN proteins or MN polypeptides comprise
5 amino acid sequences selected from the group consisting of SEQ ID NOS: 10 and 97-106. Alternatively, said MN polypeptides are selected from the group consisting of SEQ ID NOS: 10 and 97-106.

Representative MN proteins and MN polypeptides which mediate attachment of vertebrate cells in a cell adhesion assay, are specifically bound by either
10 the M75 monoclonal antibody that is secreted from the hybridoma VU-M75, which was deposited at the American Type Culture Collection under ATCC No. HB 11128, or by the MN12 monoclonal antibody that is secreted from the hybridoma MN 12.2.2, which was deposited at the American Type Culture Collection under ATCC No. HB 11647, or by both said monoclonal antibodies.

15 Another aspect of the instant invention is a method of identifying a site on an MN protein to which vertebrate cells adhere by testing a series of overlapping polypeptides from said MN protein in a cell adhesion assay with vertebrate cells, and determining that if cells adhere to a polypeptide from said series, that said polypeptide comprises a site on said MN protein to which vertebrate cells adhere.

20 Still another aspect of the instant invention is a vector comprising an expression control sequence operatively linked to a nucleic acid encoding the variable domains of a MN-specific antibody, wherein said domains are separated by a flexible linker polypeptide, and wherein said vector, when transfected into a vertebrate preneoplastic or neoplastic cell that abnormally expresses MN protein, inhibits the
25 growth of said cell. Preferably said expression control sequence comprises the MN gene promoter operatively linked to said nucleic acid. Further preferably, said flexible linker polypeptide has the amino acid sequence of SEQ ID NO: 116, and even further preferably, said MN gene promoter has the nucleotide sequence of SEQ ID NO: 27.

Another further aspect of the instant invention concerns a vector
30 comprising a nucleic acid that encodes a cytotoxic protein or cytotoxic polypeptide operatively linked to the MN gene promoter, wherein said vector, when transfected into a vertebrate preneoplastic or neoplastic cell that abnormally expresses MN protein,

inhibits the growth of said cell. In one preferred embodiment said cytotoxic protein is HSV thymidine kinase. Preferably, said vector further comprises a nucleic acid encoding a cytokine operatively linked to said MN gene promoter. In alternative and preferred embodiments, said cytokine is interferon or interleukin-2.

5 The MN gene promoter is characterized herein. The identification of the binding site for a repressor of MN transcription is disclosed. Mutational analysis indicated that the direct repeat AGGGCacAGGGC [SEQ ID NO: 143] is required for efficient repressor binding.

 Identification of the protein that binds to the repressor and modification
10 of its binding properties is another route to modulate MN expression leading to cancer therapies. Suppression of MN expression in tumor cells by over expression of a negative regulator is expected to lead to a decrease of tumor growth. A repressor complex comprising at least two subunits was found to bind to SEQ ID NO: 115 of the MN gene promoter. A repressor complex, found to be in direct contact with SEQ ID
15 NO: 115 by UV crosslinking, comprised two proteins having molecular weights of 35 and 42 kilodaltons, respectively.

Abbreviations

The following abbreviations are used herein:

20	aa	-	amino acid
	ATCC	-	American Type Culture Collection
	bp	-	base pairs
	BLV	-	bovine leukemia virus
	BSA	-	bovine serum albumin
25	BRL	-	Bethesda Research Laboratories
	CA	-	carbonic anhydrase
	CAM	-	cell adhesion molecule
	CARP	-	carbonic anhydrase related protein
	CAT	-	chloramphenicol acetyltransferase
30	Ci	-	curie
	cm	-	centimeter
	CMV	-	cytomegalovirus

	cpm	-	counts per minute
	C-terminus	-	carboxyl-terminus
	CTL	-	cytotoxic T lymphocytes
	°C	-	degrees centigrade
5	DEAE	-	diethylaminoethyl
	DMEM	-	Dulbecco modified Eagle medium
	ds	-	double-stranded
	EDTA	-	ethylenediaminetetraacetate
	EGF	-	epidermal growth factor
10	EIA	-	enzyme immunoassay
	ELISA	-	enzyme-linked immunosorbent assay
	EMSA	-	electrophoretic mobility shift assay
	F	-	fibroblasts
	FACS	-	cytofluorometric study
15	FCS	-	fetal calf serum
	FITC	-	fluorescein isothiocyanate
	FTP	-	DNase 1 footprinting analysis
	GST-MN	-	fusion protein MN glutathione S-transferase
	GVC	-	ganciclovir
20	H	-	HeLa cells
	H-E	-	haematoxylin-eosin
	HEF	-	human embryo fibroblasts
	HeLa K	-	standard type of HeLa cells
	HeLa S	-	Stanbridge's mutant HeLa D98/AH.2
25	H/F-T	-	hybrid HeLa fibroblast cells that are tumorigenic; derived from HeLa D98/AH.2
	H/F-N	-	hybrid HeLa fibroblast cells that are nontumorigenic; derived from HeLa D98/AH.2
	HPV	-	Human papilloma virus
30	HRP	-	horseradish peroxidase
	HSV	-	Herpes simplex virus
	IC	-	intracellular

	IFN	-	interferon
	IL-2	-	interleukin-2
	Inr	-	initiator
	IPTG	-	isopropyl-Beta-D-thiogalacto-pyranoside
5	kb	-	kilobase
	kbp	-	kilobase pairs
	kd or kDa	-	kilodaltons
	KS	-	keratan sulphate
	LCMV	-	lymphocytic choriomeningitis virus
10	LTR	-	long terminal repeat
	M	-	molar
	mA	-	milliampere
	MAb	-	monoclonal antibody
	MCSF	-	macrophage colony stimulating factor
15	ME	-	mercaptoethanol
	MEM	-	minimal essential medium
	min.	-	minute(s)
	mg	-	milligram
	ml	-	milliliter
20	mM	-	millimolar
	MMC	-	mitomycin C
	mmol	-	millimole
	MLV	-	murine leukemia virus
	N	-	normal concentration
25	NEG	-	negative
	ng	-	nanogram
	nm	-	nanometer
	nt	-	nucleotide
	N-terminus	-	amino-terminus
30	ODN	-	oligodeoxynucleotide
	ORF	-	open reading frame
	PA	-	Protein A

	PBS	-	phosphate buffered saline
	PCR	-	polymerase chain reaction
	PEST	-	combination of one-letter abbreviations for proline, glutamic acid, serine, threonine
5	PG	-	proteoglycan
	pI	-	isoelectric point
	PMA	-	phorbol 12-myristate 13-acetate
	POS	-	positive
	Py	-	pyrimidine
10	RACE	-	rapid amplification of cDNA ends
	RCC	-	renal cell carcinoma
	RIA	-	radioimmunoassay
	RIP	-	radioimmunoprecipitation
	RIPA	-	radioimmunoprecipitation assay
15	RNP	-	RNase protection assay
	RT-PCT	-	reverse transcription polymerase chain reaction
	SAC	-	<u>Staphylococcus aureus</u> cells
	<u>S. aureus</u>	-	<u>Staphylococcus aureus</u>
	sc	-	subcutaneous
20	SDRE	-	serum dose response element
	SDS	-	sodium dodecyl sulfate
	SDS-PAGE	-	sodium dodecyl sulfate-polyacrylamide gel electrophoresis
	SINE	-	short interspersed repeated sequence
	SP	-	signal peptide
25	SP-RIA	-	solid-phase radioimmunoassay
	SSDS	-	synthetic splice donor site
	SSH	-	subtractive suppressive PCR
	SSPE	-	NaCl (0.18 M), sodium phosphate (0.01 M), EDTA (0.001 M)
	TBE	-	Tris-borate/EDTA electrophoresis buffer
30	TC	-	tissue culture
	TCA	-	trichloroacetic acid
	TC media	-	tissue culture media

	TC	-	tissue culture
	tk	-	thymidine kinase
	TM	-	transmembrane
	TMB	-	tetramethylbenzidine
5	Tris	-	tris (hydroxymethyl) aminomethane
	μCi	-	microcurie
	μg	-	microgram
	μl	-	microliter
	μM	-	micromolar
10	VSV	-	vesicular stomatitis virus
	VV	-	vaccinia virus
	X-MLV	-	xenotropic murine leukemia virus

Cell Lines

15	AGS	—	cell line derived from a primary adenogastric carcinoma [Barranco and Townsend, <u>Cancer Res.</u> , 43: 1703 (1983) and <u>Invest. New Drugs</u> , 1: 117 (1983)]; available from the ATCC under CRL-1739;
20	BL-3	—	bovine B lymphocytes [ATCC CRL-8037; leukemia cell suspension; <u>J. Natl. Cancer Inst.</u> (Bethesda) 40: 737 (1968)];
	C33	—	a cell line derived from a human cervical carcinoma biopsy [Auersperg, N., <u>J. Nat'l. Cancer Inst.</u> (Bethesda), 32: 135-148 (1964)]; available from the ATCC under HTB-31;
25			
	C33A	—	human cervical carcinoma cells [ATCC HTB-31; <u>J. Natl. Cancer Inst.</u> (Bethesda) 32: 135 (1964)];
30	COS	—	simian cell line [Gluzman, Y., <u>Cell</u> , 23: 175 (1981)];

- 5 HeLa K – standard type of HeLa cells; aneuploid, epithelial-like cell line isolated from a human cervical adenocarcinoma [Gey et al., Cancer Res., 12: 264 (1952); Jones et al., Obstet. Gynecol., 38: 945-949 (1971)] obtained from Professor B. Korych, [Institute of Medical Microbiology and Immunology, Charles University; Prague, Czech Republic];
- 10 HeLa D98/AH.2 (also HeLa s) – Mutant HeLa clone that is hypoxanthine guanine phosphoribosyl transferase-deficient (HGPRT⁻) kindly provided by Eric J. Stanbridge [Department of Microbiology, College of Medicine, University of California, Irvine, CA (USA)] and reported in Stanbridge et al., Science, 215: 252-259 (15 Jan. 1982); parent of hybrid cells H/F-N and H/F-T, also obtained from E.J. Stanbridge;
- 15 KATO III – cell line prepared from a metastatic form of a gastric carcinoma [Sekiguichi et al., Japan J. Exp. Med., 48: 61 (1978)]; available from the ATCC under HTB-103;
- 20 NIH-3T3 – murine fibroblast cell line reported in Aaronson, Science, 237: 178 (1987);
- 25 QT35 – quail fibrosarcoma cells [ECACC: 93120832; Cell, 11: 95 (1977)];
- Raj – human Burkitt's lymphoma cell line [ATCC CCL-86; Lancet, 1: 238 (1964)];
- 30 Rat2TK⁻ – cell line (rat embryo, thymidine kinase mutant) was derived from a subclone of a 5'-bromo-deoxyuridine resistant strain of the Fischer rat fibroblast 3T3-like cell line Rat1; the cells lack

appreciable levels of nuclear thymidine kinase [Ahrens, B., Virology, 113: 408 (1981)];

- 5 SiHa – human cervical squamous carcinoma cell line [ATCC HTB-35; Friedl et al., Proc. Soc. Exp. Biol. Med., 135: 543 (1990)];
- 10 XC – cells derived from a rat rhabdomyosarcoma induced with Rous sarcoma virus-induced rat sarcoma [Svoboda, J., Natl. Cancer Center Institute Monograph No. 17, IN: "International Conference on Avian Tumor Viruses" (J.W. Beard ed.), pp. 277-298 (1964)], kindly provided by Jan Svoboda [Institute of Molecular Genetics, Czechoslovak Academy of Sciences; Prague, Czech Republic]; and
- 15 CGL1 – H/F-N hybrid cells (HeLa D98/AH.2 derivative);
- CGL2 – H/F-N hybrid cells (HeLa D98/AH.2 derivative);
- 20 CGL3 – H/F-T hybrid cells (HeLa D98/AH.2 derivative);
- CGL4 – H/F-T hybrid cells (HeLa D98/Ah.2 derivative).

Nucleotide and Amino Acid Sequence Symbols

The following symbols are used to represent nucleotides herein:

25	Base <u>Symbol</u>	<u>Meaning</u>
	A	adenine
	C	cytosine
	G	guanine
30	T	thymine
	U	uracil
	I	inosine

	M	A or C
	R	A or G
	W	A or T/U
	S	C or G
5	Y	C or T/U
	K	G or T/U
	V	A or C or G
	H	A or C or T/U
	D	A or G or T/U
10	B	C or G or T/U
	N/X	A or C or G or T/U

There are twenty main amino acids, each of which is specified by a different arrangement of three adjacent nucleotides (triplet code or codon), and which are linked together in a specific order to form a characteristic protein. A three-letter or one-letter convention is used herein to identify said amino acids, as, for example, in Figure 1 as follows:

	<u>Amino acid name</u>	<u>3 Ltr. Abbrev.</u>	<u>1 Ltr. Abbrev.</u>
20	Alanine	Ala	A
	Arginine	Arg	R
	Asparagine	Asn	N
25	Aspartic Acid	Asp	D
	Cysteine	Cys	C
	Glutamic Acid	Glu	E
	Glutamine	Gln	Q
	Glycine	Gly	G
30	Histidine	His	H
	Isoleucine	Ile	I
	Leucine	Leu	L
	Lysine	Lys	K

	Methionine	Met	M
	Phenylalanine	Phe	F
	Proline	Pro	P
	Serine	Ser	S
5	Threonine	Thr	T
	Tryptophan	Trp	W
	Tyrosine	Tyr	Y
	Valine	Val	V
	Unknown or other		X

10

BRIEF DESCRIPTION OF THE FIGURES

Figure 1A-C provides the nucleotide sequence for a MN cDNA [SEQ ID NO: 1] clone isolated as described herein. Figure 1 A-C also sets forth the predicted amino acid sequence [SEQ ID NO: 2] encoded by the cDNA.

15

Figure 2A-F provides a 10,898 bp complete genomic sequence of MN [SEQ ID NO: 5]. The base count is as follows: 2654 A; 2739 C; 2645 G; and 2859 T. The 11 exons are in general shown in capital letters, but exon 1 is considered to begin at position 3507 as determined by RNase protection assay.

20

Figure 3 is a restriction map of the full-length MN cDNA. The open reading frame is shown as an open box. The thick lines below the restriction map illustrate the sizes and positions of two overlapping cDNA clones. The horizontal arrows indicate the positions of primers R1 [SEQ ID NO: 7] and R2 [SEQ ID NO: 8] used for the 5' end RACE. Relevant restriction sites are BamHI (B), EcoRV (V), EcoRI (E), PstI (Ps), PvuII (Pv).

25

Figure 4 schematically represents the 5' MN genomic region of a MN genomic clone wherein the numbering corresponds to transcription initiation sites estimated by RACE.

30

Figure 5 provides an exon-intron map of the human MN/CA IX gene. The positions and sizes of the exons (numbered, cross-hatched boxes), Alu repeat elements (open boxes) and an LTR-related sequence (first unnumbered stippled box) are adjusted to the indicated scale. The exons corresponding to individual MN/CA IX protein domains are enclosed in dashed frames designated PG (proteoglycan-like domain), CA

(carbonic anhydrase domain), TM (transmembrane anchor) and IC (intracytoplasmic tail). Below the map, the alignment of amino acid sequences illustrates the extent of homology between the MN/CA IX protein PG region (aa 53-111) [SEQ ID NO: 50] and the human aggrecan (aa 781-839) [SEQ ID NO: 54].

5 Figure 6 is a nucleotide sequence for the proposed promoter of the human MN gene [SEQ ID NO: 27]. The nucleotides are numbered from the transcription initiation site according to RNase protection assay. Potential regulatory elements are overlined. Transcription start sites are indicated by asterisks (RNase protection) and dots (RACE) above the corresponding nucleotides. The sequence of the
10 1st exon begins under the asterisks. FTP analysis of the MN4 promoter fragment revealed 5 regions (I-V) protected at both the coding and noncoding strands, and two regions (VI and VII) protected at the coding strand but not at the noncoding strand.

 Figure 7 provides a schematic of the alignment of MN genomic clones according to their position related to the transcription initiation site. All the genomic
15 fragments except Bd3 were isolated from a lambda FIX II genomic library derived from HeLa cells. Clone Bd3 was derived from a human fetal brain library.

 Figure 8 schematically represents the MN protein structure. The abbreviations are the same as used in Figure 5. The scale indicates the number of amino acids.

20

DETAILED DESCRIPTION

 The terms "MN/CA IX" and "MN/CA9" are herein considered to be synonyms for MN. Also, the G250 antigen is considered to refer to MN
25 protein/polypeptide. [Uemura et al., J. Urol., **154** (4 Suppl.): 377 (Abstract 1475; 1997).]

 MN/CA IX was first identified in HeLa cells, derived from human carcinoma of cervix uteri, as both a plasma membrane and nuclear protein with an apparent molecular weight of 58 and 54 kilodaltons (kDA) as estimated by Western
30 blotting. It is N-glycosylated with a single 3kDa carbohydrate chain and under non-reducing conditions forms S-S-linked oligomers [Pastorekova et al., Virology, **187**: 620-626 (1992); Pastorek et al., Oncogene, **9**: 2788-2888 (1994)]. MN/CA IX is a

transmembrane protein located at the cell surface, although in some cases it has been detected in the nucleus [Zavada et al., Int. J. Cancer, 54: 268-274 (1993); Pastorekova et al., supra].

MN is manifested in HeLa cells by a twin protein, p54/58N.

- 5 Immunoblots using a monoclonal antibody reactive with p54/58N (MAb M75) revealed two bands at 54 kd and 58 kd. Those two bands may correspond to one type of protein that most probably differs by post-translational processing. Herein, the phrase "twin protein" indicates p54/58N.

- 10 Zavada et al., WO 93/18152 and/or WO 95/34650 disclose the MN cDNA sequence (SEQ ID NO: 1) shown herein in Figure 1A-1C, the MN amino acid sequence (SEQ ID NO: 2) also shown in Figure 1A-1C, and the MN genomic sequence (SEQ ID NO: 5) shown herein in Figure 2A-2F. The MN gene is organized into 11 exons and 10 introns.

- 15 The first thirty seven amino acids of the MN protein shown in Figure 1A-1C is the putative MN signal peptide [SEQ ID NO: 6]. The MN protein has an extracellular domain [amino acids (aa) 38-414 of Figure 1A-1C (SEQ ID NO: 87)], a transmembrane domain [aa 415-434 (SEQ ID NO: 52)] and an intracellular domain [aa 435-459 (SEQ ID NO: 53)]. The extracellular domain contains the proteoglycan-like domain [aa 53-111 (SEQ ID NO: 50)] and the carbonic anhydrase (CA) domain [aa 135-20 391 (SEQ ID NO: 51)].

Anticancer Drugs and Antibodies that Block Interaction of MN Protein and Receptor Molecules

- 25 MN protein is considered to be a uniquely suitable target for cancer therapy for a number of reasons including the following. (1) It is localized on the cell surface, rendering it accessible. (2) It is expressed in a high percentage of human carcinomas (e.g., uterine cervical, renal, colon, breast, esophageal, lung, head and neck carcinomas, among others), but is not normally expressed to any significant extent in
30 the normal tissues from which such carcinomas originate.

(3) It is normally expressed only in the stomach mucosa and in some epithelia of the digestive tract (epithelium of gallbladder and small intestine). An anatomic barrier thereby exists between the MN-expressing preneoplastic/neoplastic

and MN-expressing normal tissues. Drugs, including antibodies, can thus be administered which can reach tumors without interfering with MN-expressing normal tissues.

(4) MAb M75 has a high affinity and specificity to MN protein. (5) MN cDNA and MN genomic clones which encompass the protein-coding and gene regulatory sequences have been isolated. (6) MN-specific antibodies have been shown to have among the highest tumor uptakes reported in clinical studies with antitumor antibodies in solid tumors, as shown for the MN-specific chimeric antibody G250 in animal studies and in phase I clinical trials with renal carcinoma patients. [Steffens et al., J. Clin. Oncol., 15: 1529 (1997).] Also, MN-specific antibodies have low uptake in normal tissues.

Data, e.g. as presented herein, are consistent with the following theory concerning how MN protein acts in normal tissues and in preneoplastic/neoplastic tissues. In normal tissues (e.g., in stomach mucosa), MN protein is considered to be a differentiation factor. It binds with its normal receptor S (for stomach). Stomach carcinomas have been shown not to contain MN protein.

Ectopic expression of MN protein in other tissues causes malignant conversion of cells. Such ectopic expression is considered to be caused by the binding of MN protein with an alternative receptor H (for HeLa cells), coupled to a signal transduction pathway leading to malignancy. Drugs or antibodies which block the binding site of MN protein for receptor H would be expected to cause reversion of preneoplastic/neoplastic cells to normal or induce their death.

Design and Development of MN-Blocking Drugs or Antibodies

A process to design and develop MN-blocking drugs, e.g., peptides with high affinity to MN protein, or antibodies, has several steps. First, is to test for the binding of MN protein to receptors based on the cell adhesion assay described infra. That same procedure would also be used to assay for drugs blocking the MN protein binding site. In view of the alternative receptors S and H, stomach epithelial cells or revertants (containing preferentially S receptors), HeLa cells (containing the H receptor and lacking the S receptor) would be used in the cell adhesion assay.

To identify the receptor binding site of MN protein, deletion variants of MN protein lacking different domains can be used to identify region(s) responsible for interaction of MN protein with a receptor. Example 2 identifies and illustrates how to detect other binding sites on MN protein. A preferred MN binding site is considered to be closely related or identical to the epitope for MAb M75, which is located in at least 2 copies within the 6-fold tandem repeat of 6 amino acids [aa 61-96 (SEQ ID NO: 97)] in the proteoglycan-like domain of the MN protein. Smaller deletion variants can be prepared within that relevant domain, e.g., fusion proteins with only small segments of MN protein can be prepared. Also, controlled digestion of MN protein with specific proteases followed by separation of the products can be performed.

Further, peptides comprising the expected binding site can be synthesized. All of those products can be tested in cell adhesion assays, as exemplified below. [See, e.g., Pierschbacher and Ruoslahti, PNAS, **81**:5985 (1984); Ruoslahti and Pierschbacher, Science, **238**: 491.]

Molecules can be constructed to block the MN receptor binding site. For example, use of a phage display peptide library kit [as Ph.D[®]-7 Peptide 7-Mer Library Kit from New England Biolabs; Beverly, MA (USA)] as exemplified in Examples 2 and 3, can be used to find peptides with high affinity to the target molecules. Biologic activity of the identified peptides will be tested in vitro by inhibition of cell adhesion to MN protein, by effects on cell morphology and growth characteristics of MN-related tumor cells (HeLa) and of control cells. [Symington, J. Biol. Chem., **267**: 25744 (1992).] In vivo screening will be carried out in nude mice that have been injected with HeLa cells.

Peptides containing the binding site of the MN protein will be prepared [e.g. MAPs (multiple antigen peptides); Tam, J.P., PNAS (USA) **85**: 5409 (1988); Butz et al., Peptide Res., **7**: 20 (1994)]. The MAPs will be used to immunize animals to obtain antibodies (polyclonal and/or monoclonal) that recognize and block the binding site. [See, e.g., Brooks et al., Cell, **79**: 1157 (1994).] "Vaccination" would then be used to test for protection in animals. Antibodies to the MN binding site could potentially be used to block MN protein's interaction(s) with other molecules.

Computer modeling can also be used to design molecules with specific affinity to MN protein that would mediate steric inhibition between MN protein and its

receptor. A computer model of the MN binding site for the receptor will contain spatial, electrostatic, hydrophobic and other characteristics of this structure. Organic molecules complementary to the structure, that best fit into the binding site, will be designed. Inorganic molecules can also be similarly tested that could block the MN
5 binding site.

The use of oncoproteins as targets for developing new cancer therapeutics is considered conventional by those of skill in the art. [See, e.g., Mendelsohn and Lippman, "Growth Factors," pp. 114-133, IN: DeVita et al. (eds.), Cancer: Principles and Practice of Oncology (4th Ed.; Lippincott; Philadelphia, 1993).] In
10 its broadest sense, the design of blocking drugs can be based in competitive inhibition experiments. Such experiments have been used to invent drugs since the discovery of sulfonamides (competitive inhibitors of para-aminobenzoic acid, a precursor of folic acid). Also, some cytostatics are competitive inhibitors (e.g., halogenated pyrimidines, among others).

15 However, the application of such approaches to MN is new. In comparison to other tumor-related molecules (e.g. growth factors and their receptors), MN has the unique property of being differentially expressed in preneoplastic/neoplastic and normal tissues, which are separated by an anatomic barrier.

20

MN Gene – Cloning and Sequencing

Figure 1A-C provides the nucleotide sequence for a full-length MN cDNA clone isolated as described below [SEQ ID NO: 1]. Figure 2A-F provides a complete MN genomic sequence [SEQ ID NO: 5]. Figure 6 shows the nucleotide sequence for a
25 proposed MN promoter [SEQ ID NO: 27].

It is understood that because of the degeneracy of the genetic code, that is, that more than one codon will code for one amino acid [for example, the codons TTA, TTG, CTT, CTC, CTA and CTG each code for the amino acid leucine (leu)], that variations of the nucleotide sequences in, for example, SEQ ID NOS: 1 and 5 wherein
30 one codon is substituted for another, would produce a substantially equivalent protein or polypeptide according to this invention. All such variations in the nucleotide

sequences of the MN cDNA and complementary nucleic acid sequences are included within the scope of this invention.

It is further understood that the nucleotide sequences herein described and shown in Figures 1, 2 and 6, represent only the precise structures of the cDNA, genomic and promoter nucleotide sequences isolated and described herein. It is expected that slightly modified nucleotide sequences will be found or can be modified by techniques known in the art to code for substantially similar or homologous MN proteins and polypeptides, for example, those having similar epitopes, and such nucleotide sequences and proteins/ polypeptides are considered to be equivalents for the purpose of this invention. DNA or RNA having equivalent codons is considered within the scope of the invention, as are synthetic nucleic acid sequences that encode proteins/polypeptides homologous or substantially homologous to MN proteins/polypeptides, as well as those nucleic acid sequences that would hybridize to said exemplary sequences [SEQ. ID. NOS. 1, 5 and 27] under stringent conditions, or that, but for the degeneracy of the genetic code would hybridize to said cDNA nucleotide sequences under stringent hybridization conditions. Modifications and variations of nucleic acid sequences as indicated herein are considered to result in sequences that are substantially the same as the exemplary MN sequences and fragments thereof.

Stringent hybridization conditions are considered herein to conform to standard hybridization conditions understood in the art to be stringent. For example, it is generally understood that stringent conditions encompass relatively low salt and/or high temperature conditions, such as provided by 0.02 M to 0.15 M NaCl at temperatures of 50°C to 70°C. Less stringent conditions, such as, 0.15 M to 0.9 M salt at temperatures ranging from 20°C to 55°C can be made more stringent by adding increasing amounts of formamide, which serves to destabilize hybrid duplexes as does increased temperature.

Exemplary stringent hybridization conditions are described in Sambrook et al., Molecular Cloning: A Laboratory Manual, pages 1.91 and 9.47-9.51 (Second Edition, Cold Spring Harbor Laboratory Press; Cold Spring Harbor, NY; 1989); Maniatis et al., Molecular Cloning: A Laboratory Manual, pages 387-389 (Cold Spring Harbor

Laboratory; Cold Spring Harbor, NY; 1982); Tsuchiya et al., Oral Surgery, Oral Medicine, Oral Pathology, 71(6): 721-725 (June 1991).

Zavada et al., WO 95/34650 described how a partial MN cDNA clone, a full-length MN cDNA clone and MN genomic clones were isolated and sequenced.

- 5 Also, Zavada et al., Int. J. Cancer, 54: 268 (1993) describes the isolation and sequencing of a partial MN cDNA of 1397 bp in length. Briefly attempts to isolate a full-length clone from the original cDNA library failed. Therefore, the inventors performed a rapid amplification of cDNA ends (RACE) using MN-specific primers, R1 and R2 [SEQ ID NOS: 7 and 8], derived from the 5' region of the original cDNA clone.
- 10 The RACE product was inserted into pBluescript, and the entire population of recombinant plasmids was sequenced with an MN-specific primer ODN1 [SEQ ID NO: 3]. In that way, a reliable sequence at the very 5' end of the MN cDNA as shown in Figure 1 [SEQ ID NO: 1] was obtained.

- Specifically, RACE was performed using 5' RACE System [GIBCO BRL; Gaithersburg, MD (USA)] as follows. 1 μ g of mRNA (the same as above) was used as a
- 15 template for the first strand cDNA synthesis which was primed by the MN-specific antisense oligonucleotide, R1 (5'-TGGGGTTCTTGAGGATCTCCAGGAG-3') [SEQ ID NO: 7]. The first strand product was precipitated twice in the presence of ammonium acetate and a homopolymeric C tail was attached to its 3' end by TdT. Tailed cDNA
- 20 was then amplified by PCR using a nested primer, R2 (5'-CTCTAACTTCAGGGAGCCCTCTTCTT-3') [SEQ ID NO: 8] and an anchor primer that anneals to the homopolymeric tail (5'-CUACUACUACUAGGCCACGCGTCGAC TAGTACGGGI IGGGIIGGGIIG-3') [SEQ ID NO: 9]. The amplified product was
- 25 digested with BamHI and SalI restriction enzymes and cloned into pBluescript II KS plasmid. After transformation, plasmid DNA was purified from the whole population of transformed cells and used as a template for sequencing with the MN-specific primer ODN1 [SEQ ID NO: 3; a 29-mer 5' CGCCCAGTGGGTCATCTTCCCCAGAAGAG 3'].

- To study MN regulation, MN genomic clones were isolated. One MN genomic clone (Bd3) was isolated from a human cosmid library prepared from fetal
- 30 brain using both MN cDNA as a probe and the MN-specific primers derived from the 5' end of the cDNA ODN1 [SEQ ID NO: 3, supra] and ODN2 [SEQ. ID NO.: 4; 19-mer (5' GGAATCCTCCTGCATCCGG 3')]. Sequence analysis revealed that that genomic

clone covered a region upstream from a MN transcription start site and ending with the BamHI restriction site localized inside the MN cDNA. Other MN genomic clones can be similarly isolated.

Figure 7 provides a schematic of the alignment of MN genomic clones according to the transcription initiation site. Plasmids containing the A4a clone and the XE1 and XE3 subclones were deposited at the American Type Culture Collection (ATCC) on June 6, 1995, respectively under ATCC Deposit Nos. 97199, 97200, and 97198.

10 Exon-Intron Structure of Complete MN Genomic Region

The complete sequence of the overlapping clones contains 10,898 bp (SEQ ID NO: 5). Figure 5 depicts the organization of the human MN gene, showing the location of all 11 exons as well as the 2 upstream and 6 intronic Alu repeat elements. All the exons are small, ranging from 27 to 191 bp, with the exception of the first exon which is 445 bp. The intron sizes range from 89 to 1400 bp. The CA domain is encoded by exons 2-8, while the exons 1, 10 and 11 correspond respectively to the proteoglycan-like domain, the transmembrane anchor and cytoplasmic tail of the MN/CA IX protein. Table 1 below lists the splice donor and acceptor sequences that conform to consensus splice sequences including the AG-GT motif [Mount, Nucleic
20 Acids Res. 10: 459-472 (1982)].

TABLE 1

Exon-Intron Structure of the Human MN Gene

	Intron	Size	Genomic Position**	SEQ ID NO	5'splice acceptor	SEQ ID NO
5	1	445	*3507-3951	28	AGAAG gtaagt	67
	2	30	5126-5155	29	TGGAG gtgaga	68
	3	171	5349-5519	30	CAGTC gtgagg	69
	4	143	5651-5793	31	CCGAG gtgagc	70
	5	93	5883-5975	32	TGGAG gtacca	71
10	6	67	7376-7442	33	GGAAG gtcagt	72
	7	158	8777-8934	34	AGCAG gtgggc	73
	8	145	9447-9591	35	GCCAG gtacag	74
	9	27	9706-9732	36	TGCTG gtgagt	75
	10	82	10350-70431	37	CACAG gtatta	76
15	11	191	10562-10752	38	ATAAT end	
	Intron	Size	Genomic Position **	SEQ ID NO	3'splice acceptor	SEQ ID NO
	1	1174	3952-5125	39	atacag GGGAT	77
	2	193	5156-5348	40	ccccag GCGAC	78
	3	131	5520-5650	41	acgcag TGCAA	79
	4	89	5794-5882	42	tttcag ATCCA	80
	5	1400	5976-7375	43	ccccag GAGGG	81
20	6	1334	7443-8776	44	tcacag GCTCA	82
	7	512	8935-9446	45	ccctag CTCCA	83
	8	114	9592-9705	46	ctccag TCCAG	84
	9	617	9733-10349	47	tcgcag GTGACA	85
	10	130	10432-10561	48	acacag AAGGG	86

** positions are related to nt numbering in whole genomic sequence including the 5' flanking region [Figure 2A-F]

30 * number corresponds to transcription initiation site determined below by RNase protection assay

Mapping of MN Gene Transcription Initiation and Termination Sites

Zavada et al., WO 95/34650 describes the process of mapping the MN gene transcription initiation and termination sites. A RNase protection assay was used for fine mapping of the 5' end of the MN gene. The probe was a uniformly labeled 470 nucleotide copy RNA (nt -205 to +265) [SEQ ID NO: 55], which was hybridized to total RNA from MN-expressing HeLa and CGI.3 cells and analyzed on a sequencing gel. That analysis has shown that the MN gene transcription initiates at multiple sites, the 5' end of the longest MN transcript being 30 nt longer than that previously characterized by RACE.

10

Characterization of the 5' Flanking Region

The Bd3 genomic clone isolated from human fetal brain cosmid library was found to cover a region of 3.5 kb upstream from the transcription start site of the MN gene. It contains no significant coding region. Two Alu repeats are situated at positions -2587 to -2296 [SEQ ID NO: 56] and -1138 to -877 [SEQ ID NO: 57] (with respect to the transcription start determined by RNP).

Nucleotide sequence analysis of the DNA 5' to the transcription start (from nt -507) revealed no recognizable TATA box within the expected distance from the beginning of the first exon. However, the presence of potential binding sites for transcription factors suggests that this region might contain a promoter for the MN gene. There are several consensus sequences for transcription factors AP1 and AP2 as well as for other regulatory elements, including a p53 binding site [Locker and Buzard, J., DNA Sequencing and Mapping, 1: 3-11 (1990); Imagawa et al. Cell, 51: 251-260 (1987); El Deiry et al., Nat. Genet., 1: 44-49 (1992)]. Although the putative promoter region contains 59.3% C+G, it does not have additional attributes of CpG-rich islands that are typical for TATA-less promoters of housekeeping genes [Bird, Nature, 321: 209-213 (1986)]. Another class of genes lacking TATA box utilizes the initiator (Inr) element as a promoter. Many of these genes are not constitutively active, but they are rather regulated during differentiation or development. The Inr has a consensus sequence of PyPyPyCAPyPyPyPyPy [SEQ ID NO: 23] and encompasses the transcription start site [Smale and Baltimore, Cell, 57: 103-113 (1989)]. There are two such consensus

30

sequences in the MN putative promoter; however, they do not overlap the transcription start (Figure 6).

An interesting region was found in the middle of the MN gene. The region is about 1.4 kb in length [nt 4,600-6,000 of the genomic sequence; SEQ ID NO: 49] and spans from the 3' part of the 1st intron to the end of the 5th exon. The region has the character of a typical CpG-rich island, with 62.8% C + G content and 82 CpG: 131 GpC dinucleotides. Moreover, there are multiple putative binding sites for transcription factors AP2 and Sp1 [Locker and Buzard, *supra*; Briggs et al., *Science*, 234: 47-52 (1986)] concentrated in the center of this area. Particularly the 3rd intron of 131 bp in length contains three Sp1 and three AP2 consensus sequences. That data indicates the possible involvement of that region in the regulation of MN gene expression. However, functionality of that region, as well as other regulatory elements found in the proposed 5' MN promoter, remains to be determined.

15 MN Promoter

Study of the MN promoter has shown that it is TATA-less and contains regulatory sequences for AP-1, AP-2, as well as two p53 binding sites. The sequence of the 5' end of the 3.5 kb flanking region upstream of the MN gene has shown extensive homology to LTR of HERV-K endogenous retroviruses. Basal transcription activity of the promoter is very weak as proven by analyses using CAT and neo reporter genes. However, expression of the reporter genes is severalfold increased when driven from the 3.5 kb flanking region, indicating involvement of putative enhancers.

Functional characterization of the 3.5 kb MN 5' upstream region by deletion analysis lead to the identification of the [-173, +31] fragment [SEQ ID NO: 21] (also alternatively, but less preferably, the nearly identical -172, +31 fragment [SEQ ID NO: 91]) as the MN promoter. *In vitro* DNase I footprinting revealed the presence of five protected regions (PR) within the MN promoter. Detailed deletion analysis of the promoter identified PR 1 and 2 (numbered from the transcription start) as the most critical for transcriptional activity. PR4 [SEQ ID NO: 115] negatively affected transcription as its deletion led to increased promoter activity and was confirmed to function as a promoter-, position- and orientation-independent silencer element. Mutational analysis indicated that the direct repeat AGGGCacAGGGC [SEQ ID NO:

143] is required for efficient repressor binding. Two components of the repressor complex (35 and 42 kDa) were found to be in direct contact with PR4 by UV crosslinking. Increased cell density, known to induce MN expression, did not affect levels of PR4 binding in HeLa cells. Significantly reduced repressor level seems to be responsible for MN up-regulation in the case of tumorigenic CGL3 as compared to non-tumorigenic CGL1 HeLa x normal fibroblast hybrid cells.

Utility of MN Promoter as a Tumor-Specific Promoter for Gene Therapy

Being investigated is whether the MN gene promoter can be used as a tumor-specific promoter to drive the expression of a suicide gene [thymidine kinase (tk) of HSV]] and mediate the direct and bystander killing of tumor cells. HSVtk gene transferred to tumor cells converts nucleoside analogue ganciclovir (GCV) to toxic triphosphates and mediates the death of transduced and also neighboring tumor cells. The control of HSVtk by the MN gene promoter would allow its expression only in tumor cells, which are permissive for the biosynthesis of MN protein, and selectively kill such tumor cells, but not normal cells in which MN expression is repressed.

A plasmid construct in which HSVtk was cloned downstream of the MN promoter region Bd3, containing both proximal and distant regulatory elements of MN, was prepared. That plasmid pMN-HSVtk was transfected to Rat2TK- cells and C33 human cervical carcinoma cells using calcium phosphate precipitation and lipofection, respectively. Transfectants were tested for expression of HSVtk and GVC sensitivity. Analysis of the transfectants has shown the remarkable cytotoxic in vitro effect of GVC even in low concentrations (up to 95% of cells killed).

Polyclonal rabbit antiserum against HSVtk, using fusion protein with GST in pGEX-3X, has been prepared to immunodetect HSVtk synthesized in transfected cells. This model system is being studied to estimate the bystander effect, the inhibition of cloning efficiency and invasiveness of transduced and GVC-treated cells to collagen matrices. A recombinant retroviral vector with the MN promoter-driven HSVtk is to be prepared to test its in vivo efficacy using an animal model (e.g., SCID-mouse).

MN Promoter Analysis

Since the MN promoter is weak, a classical approach to study it would be limited due to the relatively low efficiency of transient transfections (up to 10%).

Therefore, stable clonal cell lines expressing constructs containing the MN promoter fused to the CAT gene were prepared. In such clonal lines, 100% of the cells express the CAT gene driven from the MN promoter, and thus, the activity of the promoter is detectable easier than in transient experiments. Also, the promoter activity can be analysed repeatedly in the same cells under different conditions or treated by different factors and drugs. This approach allows for the study of the mechanisms underlying MN regulation at the level of transcription initiation.

Several types of transfections with promoter constructs linked to a reporter CAT gene (calcium precipitation, DEAE dextran combined with DMSO shock and/or chloroquine, as well as electroporation), different methods of CAT activity assay (scintillation method, thin layer chromatography) and several recipient cell lines differing in the level of MN expression and in transfection efficiency (HeLa, SiHa, CGL3, KATO III, Rat2TK⁻ and C33 cells). Activity of the MN promoter was detected preferably by the electroporation of CGL3 cells and thin layer chromatography. Further preferably, C33 cells cotransfected with MN promoter-CAT constructs and pSV2neo were used.

1. To detect basal activity of the MN promoter and to estimate the position of the core promoter, expression of the CAT gene from constructs pMN1 to pMN7 after transfection to CGL3 cells was analyzed. Plasmids with progressive 5' deletions were transfected into CGL3 cells and activity was analyzed by CAT assay. [8 μ g of DNA was used for transfection in all cases except pBLV-LTR (2 μ g).]

Only very weak CAT activity was detected in cells transfected by pMN1 and pMN2 (containing respectively 933 bp and 600 bp of the promoter sequence). A little higher activity was exhibited with the constructs pMN3, pMN4 and pMN6 (containing respectively 446 bp, 243 bp and 58 bp of the promoter). A slight peak of activity was obtained with pMN5 (starting at position -172 with respect to the transcription start.) Thus, the function of the MN core promoter can be assigned to a region of approximately 500 bp immediately upstream from the MN transcription initiation site.

Interestingly, the activity of the large Bd3 region (covering 3.5 kbp upstream of the transcription start) was severalfold higher than the activity of the core promoter. However, its level was still much lower than that exhibited by a positive control, i.e., BLV-LTR transactivated by Tax, and even lower than the activity of BLV-LTR without transactivation. That the activity of Bd3 was elevated in comparison to the core promoter suggests the presence of some regulatory elements. Such elements are most probably situated in the sequence between pMN1 and Bd3 (i.e. from -1 kbp to -3.5 kbp) [SEQ ID NO: 58]. The cloning and transfection of several deletion versions of Bd3 covering the indicated region can be used to determine the location of the putative regulatory elements.

Similar results were obtained from transfecting KATO III cells with Bd3 and pMN4. The transfected cells expressed a lower level of MN than the CGL3 cells. Accordingly, the activity of the MN promoter was found to be lower than in CGL3 cells.

2. In a parallel approach to study the MN promoter, an analysis based on G418 selection of cells transfected by plasmids containing the promoter of interest cloned upstream from the neo gene was made. This approach is suitable to study weak promoters, since its sensitivity is much higher than that of a standard CAT assay. The principle underlying the method is as follows: an active promoter drives expression of the neo gene which protects transfected cells from the toxic effect of G418, whereas an inactive promoter results in no neo product being made and the cells transfected thereby die upon the action of G418. Therefore, the activity of the promoter can be estimated according to the number of cell colonies obtained after two weeks of selection with G418. Three constructs were used in the initial experiments – pMN1neo, pMN4neo and pMN7neo. As pMN7neo contains only 30 bp upstream of the transcription start site, it was considered a negative control. As a positive control, pSV2neo with a promoter derived from SV40 was used. Rat2TK⁻ cells were chosen as the recipient cells, since they are transfectable with high efficiency by the calcium precipitation method.

After transfection, the cells were subjected to two weeks of selection. Then the medium was removed, the cells were rinsed with PBS, and the colonies were rendered visible by staining with methylene blue. The results obtained from three

independent experiments corroborated the data from the CAT assays. The promoter construct pMN4neo exhibited higher transcriptional activity than pMN1neo. However, the difference between the positive control and pMN4neo was not so striking as in the CAT assay. That may have been due to both lower promoter activity of pSV2neo compared to Tax-transactivated pBLV-LTR and to different conditions for cell growth after transfection. From that point of view, stable transfection is probably more advantageous for MN expression, since the cells grow in colonies with close cell to cell contact, and the experiment lasts much longer, providing a better opportunity to detect promoter activity.

3. Stable transfectants expressing MN promoter-CAT chimeric genes were prepared by the cotransfection of relevant plasmids with pSV2neo. As recipient cells, HeLa cells were used first. However, no clones expressing the promoter-CAT constructs were obtained. That negative result was probably caused by homologous recombination of the transfected genomic region of MN (e.g. the promoter) with the corresponding endogenous sequence. On the basis of that experience, C33 cells derived from a HPV-negative cervical carcinoma were used. C33 cells do not express MN, since during the process of tumorigenesis, they lost genetic material including chromosomal region 9p which contains the MN gene. In these experiments, the absence of the MN gene may represent an advantage as the possibility of homologous recombinations is avoided.

C33 Cells Transfected with MN Promoter-CAT Constructs

C33 cells expressing the CAT gene under MN promoter regions Bd3 (-3500/+31) [SEQ ID NO: 90] and MN5 (-172/+31) [SEQ ID NO: 91] were used for initial experiments to analyze the influence of cell density on the transcriptional activity of the MN promoter. The results indicated that signals generated after cells come into close contact activate transcription of the CAT protein from the MN promoter in proportion to the density of the cell culture. Interestingly, the data indicated that the MN protein is not required for this phase of signal transduction, since the influence of density is clearly demonstrated in MN-negative C33 cells. Rather, it appears that MN protein acts as an effector molecule produced in dense cells in order to perform a certain biological function (i.e., to perturb contact inhibition). Also interestingly, the

MN promoter activity is detectable even in very sparse cell cultures suggesting that MN is expressed at a very low level also in sparse subconfluent culture.

Deletion Variants. Deletion variants of the Bd3-CAT promoter construct were then prepared. The constructs were cotransfected with pSV2neo into C33 cervical cells. After selection with G418, the whole population of stably transfected cells were subjected to CAT ELISA analysis. Expression of the deletion constructs resulted in the synthesis of similar levels of CAT protein to that obtained with the Bd3-CAT construct. On the basis of that preliminary data, the inventors proposed that sequences stimulating transcription of MN are located between -3506 and -3375 bp [SEQ ID NO: 92] upstream from the transcription start. That is the sequence exhibiting homology to HERV-K LTR.

However, transient transfection studies in CGL3 cells repeatedly revealed that the LTR region is not required for the enhancement of basal MN promoter activity. Further, results obtained in CGL3 cells indicate that the activating element is localized in the region from -933 to -2179 [SEQ ID NO: 110] with respect to transcription initiation site (the position of the region having been deduced from overlapping sequences in the Bd3 deletion mutants).

Interaction of Nuclear Proteins with MN Promoter Sequences

In order to identify transcription factors binding to the MN promoter and potentially regulating its activity, a series of analyses using an electrophoretic mobility shift assay (EMSA) and DNase I footprinting analysis (FTP) were performed.

EMSA

In the EMSA, purified promoter fragments MN4 (-243/+ 31) [SEQ ID NO: 93], MN5 (-172/+ 31) [SEQ ID NO: 91], MN6 (-58/+ 31) [SEQ ID NO: 94] and pMN7 (-30/+ 31) [SEQ ID NO: 95], labeled at the 3' ends by Klenow enzyme, were allowed to interact with proteins in nuclear extracts prepared from CGL1 and CGL3 cells. [40 μ g of nuclear proteins were incubated with 30,000 cpm end-labeled DNA fragments in the presence of 2 μ g poly(dIdC).] DNA-protein complexes were analysed by PAGE (native 6%), where the complexes created extra bands that migrated more slowly than

the free DNA fragments, due to the shift in mobility which is dependent on the moiety of bound protein.

The EMSA of the MN4 and MN5 promoter fragments revealed several DNA-protein complexes; however, the binding patterns obtained respectively with CGL1 and CGL3 nuclear extracts were not identical. There is a single CGL-1 specific complex.

The EMSA of the MN6 promoter fragment resulted in the formation of three identical complexes with both CGL1 and CGL3 nuclear extracts, whereas the MN7 promoter fragment did not bind any nuclear proteins.

The EMSA results indicated that the CGL1 nuclear extract contains a specific factor, which could participate in the negative regulation of MN expression in CGL1 cells. Since the specific DNA-protein complex is formed with MN4 (-243/+31) [SEQ. ID NO.: 93] and MN5 (-172/+31) [SEQ. ID NO.: 91] promoter fragments, but not with MN6 (-58/+31) [SEQ ID NO: 94], it appears that the binding site of the protein component of that specific complex is located between -173 and -58 bp [SEQ. ID NO.: 96] with respect to transcription initiation.

The next step was a series of EMSA analyses using double stranded (ds) oligonucleotides designed according to the protected regions in FTP analysis. A ds oligonucleotide derived from the protected region PR2 [covering the sequence from -72 to -56 bp (SEQ ID NO: 111)] of the MN promoter provided confirmation of the binding of the AP-1 transcription factor in competitive EMSA using commercial ds oligonucleotides representing the binding site for AP-1.

EMSA of ds oligonucleotides derived from the protected regions of PR1 [-46 to -24 bp (SEQ ID NO: 112)], PR2 [-72 to -56 bp (SEQ ID NO: 111)], PR3 [-102 to -85 (SEQ ID NO: 113)] and PR5 [-163 to -144 (SEQ ID NO: 114)] did not reveal any differences in the binding pattern of nuclear proteins extracted from CGL1 and CGL3 cells, indicating that those regions do not bind crucial transcription factors which control activation of the MN gene in CGL3, or its negative regulation in CGL1. However, EMSA of ds oligonucleotides from the protected region PR4 [-133 to -108; SEQ ID NO: 115] repeatedly showed remarkable quantitative differences between binding of CGL1 and CGL3 nuclear proteins. CGL1 nuclear proteins formed a substantially higher amount of DNA-protein complexes, indicating that the PR4 region

contains a binding site for specific transcription factor(s) that may represent a negative regulator of MN gene transcription in CGL1 cells. That fact is in accord with the previous EMSA data which showed CGL-1 specific DNA-protein complex with the promoter fragments pMN4 (-243/+ 31; SEQ ID NO: 93) and pMN5 (-172/+ 31; SEQ ID NO: 91), but not with pMN6 (-58/+ 31; SEQ ID NO: 94).

To identify the protein involved or the formation of a specific complex with the MN promoter in the PR4 region, relevant ds oligonucleotides covalently bound to magnetic beads will be used to purify the corresponding transcription factor. Alternatively the ONE Hybrid System® [Clontech (Palo Alto, CA (USA))] will be used to search for and clone transcription factors involved in regulation of the analysed promoter region. A cDNA library from HeLa cells will be used for that investigation.

FTP

To determine the precise location of cis regulatory elements that participate in the transcriptional regulation of the MN gene, FTP was used. Proteins in nuclear extracts prepared respectively from CGL1 and CGL3 cells were allowed to interact with a purified ds DNA fragment of the MN promoter (MN4, -243/+ 31) [SEQ ID NO: 93] which was labeled at the 5' end of one strand. [MN4 fragments were labeled either at Xho1 site (-243/+ 31*) or at Xba1 site (*-243/+ 31).] The DNA-protein complex was then subjected to DNase I attack, which causes the DNA chain to break at certain bases if they are not in contact with proteins. [A control used BSA instead of DNase.] Examination of the band pattern of the denatured DNA after gel electrophoresis [8% denaturing gel] indicates which of the bases on the labeled strand were protected by protein.

FTP analysis of the MN4 promoter fragment revealed 5 regions (I-V) protected at both the coding and noncoding strand, as well as two regions (VI and VII) protected at the coding strand but not at the noncoding strand. Figure 6 indicates the general regions on the MN promoter that were protected.

The sequences of the identified protected regions (PR) were subjected to computer analysis using the SIGNALSCAN program to see if they corresponded to known consensus sequences for transcription factors. The data obtained by that computer analyses are as follows:

- PR I - coding strand - AP-2, p53, GAL4 noncoding strand - JCV-repeated
- PR II - coding strand - AP-1, CGN4 noncoding strand - TCF-1, dFRA, CGN4
- PR III - coding strand - no known consensus sequence, only partial overlap of AP1 noncoding strand - 2 TCF-1 sites
- 5 PR IV - coding strand - TCF-1, ADR-1 noncoding strand - CTCF, LF-A1, LBP-1
- PR V - coding strand - no known consensus motif noncoding strand - JCV repeated
- PR VI - coding strand - no known consensus motif noncoding strand - T antigen of SV 40, GAL4
- 10 PR VII - coding strand - NF-uE4, U2snRNA.2 noncoding strand - AP-2, IgHC.12, MyoD.

In contrast to EMSA, the FTP analysis did not find any differences between CGL1 and CGL3 nuclear extracts. However, the presence of specific DNA-protein interactions detected in the CGL1 nuclear extracts by EMSA could have resulted from the binding of additional protein to form DNA protein-protein complex. If that specific protein did not contact the DNA sequence directly, its presence would not be detectable by FTP.

20 EMSA Supershift Analysis

The results of the FTP suggests that transcription factors AP-1, AP-2 as well as tumor suppressor protein p53 are potentially involved in the regulation of MN expression. To confirm binding of those particular proteins to the MN promoter, a supershift analysis using antibodies specific for those proteins was performed. For this analysis, DNA-protein complexes prepared as described for EMSA were allowed to interact with MAbs or polyclonal antibodies specific for proteins potentially included in the complex. The binding of antibody to the corresponding protein results in an additional shift (supershift) in mobility of the DNA-protein-antibody complex which is PAGE visualized as an additional, more slowly migrating band.

30 By this method, the binding of AP-2 to the MN promoter was confirmed. However, this method did not evidence binding of the AP-1 transcription factor. It is

possible that MN protein binds AP-1-related protein, which is antigenically different from the AP-1 recognized by the antibodies used in this assay.

Also of high interest is the possible binding of the p53 tumor suppressor protein to the MN promoter. It is well known that wt p53 functions as a transcription factor, which activates expression of growth-restricting genes and down-modulates, directly or indirectly, the expression of genes that are required for ongoing cell proliferation. Transient co-transfection experiments using the pMN4-CAT promoter construct in combination with wt p53 cDNA and mut p53 cDNA, respectively, suggested that wt p53, but not mut p53, negatively regulates expression of MN. In addition, one of two p53-binding sites in the MN promoter is protected in FTP analysis (Figure 6), indicating that it binds to the corresponding protein. Therefore, supershift analysis to prove that p53 binds to the MN promoter with two p53-specific antibodies, e.g. Mabs 421 and DO-1 [the latter kindly provided by Dr. Vojtesek from Masaryk Memorial Cancer Institute in Brno, Czech Republic] are to be performed with appropriate nuclear extracts, e.g. from MCF-7 breast carcinoma cells which express wt p53 at a sufficient level.

Regulation of MN Expression and MN Promoter

MN appears to be a novel regulatory protein that is directly involved in the control of cell proliferation and in cellular transformation. In HeLa cells, the expression of MN is positively regulated by cell density. Its level is increased by persistent infection with LCMV. In hybrid cells between HeLa and normal fibroblasts, MN expression correlates with tumorigenicity. The fact that MN is not present in nontumorigenic hybrid cells (CGL1), but is expressed in a tumorigenic segregant lacking chromosome 11, indicates that MN is negatively regulated by a putative suppressor in chromosome 11.

Evidence supporting the regulatory role of MN protein was found in the generation of stable transfectants of NIH 3T3 cells that constitutively express MN protein. As a consequence of MN expression, the NIH 3T3 cells acquired features associated with a transformed phenotype: altered morphology, increased saturation density, proliferative advantage in serum-reduced media, enhanced DNA synthesis and capacity for anchorage-independent growth. Further, flow cytometric analyses of

asynchronous cell populations indicated that the expression of MN protein leads to accelerated progression of cells through G1 phase, reduction of cell size and the loss of capacity for growth arrest under inappropriate conditions. Also, MN expressing cells display a decreased sensitivity to the DNA damaging drug mitomycin C.

5 Nontumorigenic human cells, CGL1 cells, were also transfected with the full-length MN cDNA. The same pSG5C-MN construct in combination with pSV2neo plasmid as used to transfect the NIH 3T3 cells was used. Out of 15 MN-positive clones (tested by SP-RIA and Western blotting), 3 were chosen for further analysis. Two MN-negative clones isolated from CGL1 cells transfected with empty plasmid were added as
10 controls. Initial analysis indicates that the morphology and growth habits of MN-transfected CGL1 cells are not changed dramatically, but their proliferation rate and plating efficiency is increased.

MN Promoter – Sense/Antisense Constructs

15 When the promoter region from the MN genomic clone, isolated as described above, was linked to MN cDNA and transfected into CGL1 hybrid cells, expression of MN protein was detectable immediately after selection. However, then it gradually ceased, indicating thus an action of a feedback regulator. The putative regulatory element appeared to be acting via the MN promoter, because when the full-
20 length cDNA (not containing the promoter) was used for transfection, no similar effect was observed.

 An "antisense" MN cDNA/MN promoter construct was used to transfect CGL3 cells. The effect was the opposite of that of the CGL1 cells transfected with the "sense" construct. Whereas the transfected CGL1 cells formed colonies several times
25 larger than the control CGL1, the transfected CGL3 cells formed colonies much smaller than the control CGL3 cells. The same result was obtained by antisense MN cDNA transfection in SiHa and HeLa cells.

 For those experiments, the part of the promoter region that was linked to the MN cDNA through a BamHI site was derived from a NcoI - BamHI fragment of the
30 MN genomic clone [Bd3] and represents a region a few hundred bp upstream from the transcription initiation site. After the ligation, the joint DNA was inserted into a pBK-CMV expression vector [Stratagene]. The required orientation of the inserted sequence

was ensured by directional cloning and subsequently verified by restriction analysis. The transfection procedure was the same as used in transfecting the NIH 3T3 cells, but co-transfection with the pSV2neo plasmid was not necessary since the neo selection marker was already included in the pBK-CMV vector.

5 After two weeks of selection in a medium containing G418, remarkable differences between the numbers and sizes of the colonies grown were evident as noted above. Immediately following the selection and cloning, the MN-transfected CGL1 and CGL3 cells were tested by SP-RIA for expression and repression of MN, respectively. The isolated transfected CGL1 clones were MN positive (although the
10 level was lower than obtained with the full-length cDNA), whereas MN protein was almost absent from the transfected CGL3 clones. However, in subsequent passages, the expression of MN in transfected CGL1 cells started to cease, and was then blocked perhaps evidencing a control feedback mechanism.

 As a result of the very much lowered proliferation of the transfected
15 CGL3 cells, it was difficult to expand the majority of cloned cells (according to SP-RIA, those with the lowest levels of MN), and they were lost during passaging. However, some clones overcame that problem and again expressed MN. It is possible that once those cells reached a higher quantity, that the level of endogenously produced MN mRNA increased over the amount of ectopically expressed antisense mRNA.

20

Identification of Specific Transcription Factors Involved in Control of MN Expression

 Control of MN expression at the transcription level involves regulatory
25 elements of the MN promoter. Those elements bind transcription factors that are responsible for MN activation in tumor cells and/or repression in normal cells. The identification and isolation of those specific transcription factors and an understanding of how they regulate MN expression could result in their therapeutic utility in modulating MN expression.

30 EMSA experiments indicate the existence of an MN gene repressor. Using the One Hybrid System® [Clontech (Palo Alto, CA); an in vivo yeast genetic assay for isolating genes encoding proteins that bind to a target, cis-acting regulatory element or any other short DNA-binding sequence; Fields and Song, Nature, 340: 245 (1989);

Wu et al., EMBO J., 13: 4823 (1994)] and subtractive suppressive PCR (SSH). SSH allows the cloning of genes that are differentially expressed under conditions which are known to up or down regulate MN expression such as density versus sparsity of HeLa cells, and suspension versus adherent HeLa cells.

5 In experiments with HPV immobilized cervical cells (HCE 16/3), it was found that the regulation of MN expression differs from that in fully transformed carcinoma cells. For example, glucocorticoid hormones, which activate HPV transcription, negatively regulate MN expression in HCE, but stimulate MN in HeLa and SiHa. Further keratinocyte growth factors, which down regulates transcription of HPV
10 oncogenes, stimulates MN expression in suspension HCE but not in adherent cells.

EGF and insulin are involved in the activation of MN expression in both immortalized and carcinoma cells. All the noted facts can be used in the search for MN-specific transcription factors and in the modulation of MN expression for therapeutic purposes.

15

Deduced Amino Acid Sequence

The ORF of the MN cDNA shown in Figure 1 has the coding capacity for a 459 amino acid protein with a calculated molecular weight of 49.7 kd. The overall amino acid composition of the MN/CA IX protein is rather acidic, and predicted to have
20 a pi of 4.3. Analysis of native MN/CA IX protein from CGL3 cells by two-dimensional electrophoresis followed by immunoblotting has shown that in agreement with computer prediction, the MN/CA IX is an acidic protein existing in several isoelectric forms with pIs ranging from 4.7 to 6.3.

As assessed by amino acid sequence analysis, the deduced primary
25 structure of the MN protein can be divided into four distinct regions. The initial hydrophobic region of 37 amino acids (aa) corresponds to a signal peptide. The mature protein has an N-terminal or extracellular part of 377 amino acids [aa 38-414 (SEQ ID NO: 87)], a hydrophobic transmembrane segment of 20 amino acids [aa 415-434 (SEQ ID NO: 52)] and a C-terminal region of 25 amino acids [aa 435-459 (SEQ ID NO: 53)].

30 The extracellular part is composed of two distinct domains: (1) a proteoglycan-like domain [aa 53-111 (SEQ ID NO: 50)]; and (2) a CA domain, located

close to the plasma membrane [aa 135-391 (SEQ ID NO: 51)]. [The amino acid numbers are keyed to those of Figure 1.]

More detailed insight into MN protein primary structure disclosed the presence of several consensus sequences. One potential N-glycosylation site was found at position 346 of Figure 1. That feature, together with a predicted membrane-spanning region are consistent with the results, in which MN was shown to be an N-glycosylated protein localized in the plasma membrane. MN protein sequence deduced from cDNA was also found to contain seven S/TPXX sequence elements [SEQ ID NOS: 25 AND 26] (one of them is in the signal peptide) defined by Suzuki, J. Mol. Biol., 207: 61-84 (1989) as motifs frequently found in gene regulatory proteins. However, only two of them are composed of the suggested consensus amino acids.

Experiments have shown that the MN protein is able to bind zinc cations, as shown by affinity chromatography using Zn-charged chelating sepharose. MN protein immunoprecipitated from HeLa cells by Mab M75 was found to have weak catalytic activity of CA. The CA-like domain of MN has a structural predisposition to serve as a binding site for small soluble domains. Thus, MN protein could mediate some kind of signal transduction.

MN protein from LCMV-infected HeLa cells was shown by using DNA cellulose affinity chromatography to bind to immobilized double-stranded salmon sperm DNA. The binding activity required both the presence of zinc cations and the absence of a reducing agent in the binding buffer.

CA Domain Required for Anchorage Independence But for Increased Proliferation of Transfected NIH 3T3 Fibroblasts

In transfected NIH 3T3 fibroblasts, MN protein induces morphologic transformation, increased proliferation and anchorage independence. The consequences of constitutive expression of two MN-truncated variants in NIH 3T3 cells were studied. It was found that the proteoglycan-like region is sufficient for the morphological alteration of transfected cells and displays the growth-promoting activity presumably related to perturbation of contact inhibition.

The CA domain is essential for induction of anchorage independence, whereas the TM anchor and IC tail are dispensable for that biological effect. The MN protein is also capable of causing plasma membrane ruffling in the transfected cells and appears to participate in their attachment to the solid support. The data evince the involvement of MN in the regulation of cell proliferation, adhesion and intercellular communication.

Sequence Similarities

Computer analysis of the MN cDNA sequence was carried out using DNASIS and PROSIS (Pharmacia Software packages). GenBank, EMBL, Protein Identification Resource and SWISS-PROT databases were searched for all possible sequence similarities. In addition, a search for proteins sharing sequence similarities with MN was performed in the MIPS databank with the FastA program [Pearson and Lipman, PNAS (USA), 85: 2444 (1988)].

The proteoglycan-like domain [aa 53-111 (SEQ ID NO: 50)], which is between the signal peptide and the CA domain, shows significant homology (38% identity and 44% positivity) with a keratan sulphate attachment domain of a human large aggregating proteoglycan aggrecan [Doege et al., J. Biol. Chem., 266: 894-902 (1991)].

The CA domain [aa 135-391 (SEQ ID NO: 51)] is spread over 265 aa and shows 38.9% amino acid identity with the human CA VI isoenzyme [Aldred et al., Biochemistry, 30: 569-575 (1991)]. The homology between MN/CA IX and other isoenzymes is as follows: 35.2% with CA II in a 261 aa overlap [Montgomery et al., Nucl. Acids. Res., 15: 4687 (1987)], 31.8% with CA I in a 261 aa overlap [Barlow et al., Nucl. Acids Res., 15: 2386 (1987)], 31.6% with CA IV in a 266 aa overlap [Okuyama et al., PNAS (USA) 89: 1315-1319 (1992)], and 30.5% with CA III in a 259 aa overlap [Lloyd et al., Genes. Dev., 1: 594-602 (1987)].

In addition to the CA domain, MN/CA IX has acquired both N-terminal and C-terminal extensions that are unrelated to the other CA isoenzymes. The amino acid sequence of the C-terminal part, consisting of the transmembrane anchor and the intracytoplasmic tail, shows no significant homology to any known protein sequence.

The MN gene was clearly found to be a novel sequence derived from the human genome. The overall sequence homology between the cDNA MN sequence and cDNA sequences encoding different CA isoenzymes is in a homology range of 48-50% which is considered by ones in the art to be low. Therefore, the MN cDNA sequence is not closely related to any CA cDNA sequences.

Only very closely related nt sequences having a homology of at least 80-90% would hybridize to each other under stringent conditions. A sequence comparison of the MN cDNA sequence shown in Figure 1 and a corresponding cDNA of the human carbonic anhydrase II (CA II) showed that there are no stretches of identity between the two sequences that would be long enough to allow for a segment of the CA II cDNA sequence having 25 or more nucleotides to hybridize under stringent hybridization conditions to the MN cDNA or vice versa.

A search for nt sequences related to MN gene in the EMBL Data Library did not reveal any specific homology except for 6 complete and 2 partial Alu-type repeats with homology to Alu sequences ranging from 69.8% to 91% [Jurka and Milosavljevic, *J. Mol. Evol.* 32: 105-121 (1991)]. Also a 222 bp sequence proximal to the 5' end of the genomic region is shown to be closely homologous to a region of the HERV-K LTR.

In general, nucleotide sequences that are not in the Alu or LTR-like regions, of preferably 25 bases or more, or still more preferably of 50 bases or more, can be routinely tested and screened and found to hybridize under stringent conditions to only MN nucleotide sequences. Further, not all homologies within the Alu-like MN genomic sequences are so close to Alu repeats as to give a hybridization signal under stringent hybridization conditions. The percent of homology between MN Alu-like regions and a standard Alu-J sequence are indicated as follows:

<u>Region of Homology within</u> <u>MN Genomic Sequence</u> <u>[SEQ ID NO: 5;</u> <u>Figure 2A-F]</u>	<u>SEQ.</u> <u>ID.</u> <u>NOS.</u>	<u>% Homology to</u> <u>Entire Alu-J</u> <u>Sequence</u>
921-1212	59	89.1%
2370-2631	60	78.6%
4587-4880	61	90.1%

	6463-6738	62	85.4%
	7651-7939	63	91.0%
	9020-9317	64	69.8%
			<u>% Homology to</u> <u>One Half of</u> <u>Alu-J Sequence</u>
5	8301-8405	65	88.8%
	10040-10122	66	73.2%.

MN Proteins and/or Polypeptides

10 The phrase "MN proteins and/or polypeptides" (MN proteins/polypeptides) is herein defined to mean proteins and/or polypeptides encoded by an MN gene or fragments thereof. An exemplary and preferred MN protein according to this invention has the deduced amino acid sequence shown in Figure 1. Preferred MN proteins/polypeptides are those proteins and/or polypeptides that have
15 substantial homology with the MN protein shown in Figure 1. For example, such substantially homologous MN proteins/ polypeptides are those that are reactive with the MN-specific antibodies of this invention, preferably the Mabs M75, MN12, MN9 and MN7 or their equivalents.

 A "polypeptide" or "peptide" is a chain of amino acids covalently bound
20 by peptide linkages and is herein considered to be composed of 50 or less amino acids. A "protein" is herein defined to be a polypeptide composed of more than 50 amino acids. The term polypeptide encompasses the terms peptide and oligopeptide.

 MN proteins exhibit several interesting features: cell membrane localization, cell density dependent expression in HeLa cells, correlation with the tumorigenic phenotype of HeLa x fibroblast somatic cell hybrids, and expression in
25 several human carcinomas among other tissues. MN protein can be found directly in tumor tissue sections but not in general in counterpart normal tissues (exceptions noted infra as in normal gastric mucosa and gallbladder tissues). MN is also expressed sometimes in morphologically normal appearing areas of tissue specimens exhibiting
30 dysplasia and/or malignancy. Taken together, these features suggest a possible

involvement of MN in the regulation of cell proliferation, differentiation and/or transformation.

It can be appreciated that a protein or polypeptide produced by a neoplastic cell in vivo could be altered in sequence from that produced by a tumor cell in cell culture or by a transformed cell. Thus, MN proteins and/or polypeptides which have varying amino acid sequences including without limitation, amino acid substitutions, extensions, deletions, truncations and combinations thereof, fall within the scope of this invention. It can also be appreciated that a protein extant within body fluids is subject to degradative processes, such as, proteolytic processes; thus, MN proteins that are significantly truncated and MN polypeptides may be found in body fluids, such as, sera. The phrase "MN antigen" is used herein to encompass MN proteins and/or polypeptides.

It will further be appreciated that the amino acid sequence of MN proteins and polypeptides can be modified by genetic techniques. One or more amino acids can be deleted or substituted. Such amino acid changes may not cause any measurable change in the biological activity of the protein or polypeptide and result in proteins or polypeptides which are within the scope of this invention, as well as, MN muteins.

The MN proteins and polypeptides of this invention can be prepared in a variety of ways according to this invention, for example, recombinantly, synthetically or otherwise biologically, that is, by cleaving longer proteins and polypeptides enzymatically and/or chemically. A preferred method to prepare MN proteins is by a recombinant means. Particularly preferred methods of recombinantly producing MN proteins are described below for the GST-MN, MN 20-19, MN-Fc and MN-PA proteins.

25

Recombinant Production of MN Proteins and Polypeptides

A representative method to prepare the MN proteins shown in Figure 1 or fragments thereof would be to insert the full-length or an appropriate fragment of MN cDNA into an appropriate expression vector as exemplified below. In Zavada et al., WO 93/18152, supra, production of a fusion protein GEX-3X-MN (now termed GST-MN) using the partial cDNA clone (described above) in the vector pGEX-3X (Pharmacia)

30

Zavada et al., WO 95/34650 describes the recombinant production of both a glycosylated MN protein expressed from insect cells and a nonglycosylated MN protein expressed from *E. coli* using the expression plasmid pEt-22b [Novagen Inc.; Madison, WI (USA)]. Recombinant baculovirus express vectors were used to infect insect cells. The glycosylated MN 20-19 protein was recombinantly produced in baculovirus-infected sf9 cells [Clontech; Palo Alto, CA (USA)]. The MN 20-19 protein misses the putative signal peptide (aas 1-37) of SEQ ID NO: 6 (Figure 1), has a methionine (Met) at the N-terminus for expression, and a Leu-Glu-His-His-His-His-His [SEQ. ID NO.: 22] added to the C-terminus for purification.

Primer #20:N-terminus

Primer #19:C-terminus

The SEQ ID NOS: 17 and 18 primers were used to amplify the MN coding sequence present in the GEX-3X-MN vector using standard PCR techniques. The resulting PCR product (termed MN 20-19) was electrophoresed on a 0.5% agarose/1X TBE gel; the 1.3

kb band was excised; and the DNA recovered using the Gene Clean II kit according to the manufacturer's instructions [Bio101; LaJolla, CA (USA)].

Identification of MN Protein Partner(s)

5 A search for protein(s) interacting with MN was initiated using expression cloning of the corresponding cDNA(s) and a MN-Fc fusion protein as a probe. The chimerical MN-Fc cDNA was constructed in pSG5C vector by substitution of MN cDNA sequences encoding both the transmembrane anchor and the intracellular tail of MN protein with the cDNA encoding Fc fragment of the mouse IgG. The Fc fragment
10 cDNA was prepared by RT-PCR from the mouse hybridoma producing IgG2a antibody.

The chimerical MN-Fc cDNA was expressed by transient transfection in COS cells. COS cells were transfected using leptofection. Recombinant MN-Fc protein was released to TC medium of the transfected cells (due to the lack of the transmembrane region), purified by affinity chromatography on a Protein A Sepharose
15 and used for further experiments.

Protein extracts from mock-transfected cells and the cells transfected with pSG5C-MN-Fc were analysed by immunoblotting using the M75 MAb, Sw α M-Px and ECL detection® [ECL® – enhanced chemoluminescent system to detect phosphorylated tyrosine residues; Amersham; Arlington, Hts., IL (USA)]. The size of MN-Fc protein
20 expressed from the pSG5C vector corresponds to its computer predicted molecular weight.

³⁵S-labeled MN-Fc protein was employed in cell surface binding assay. It was found to bind to several mammalian cells, e.g., HeLa, Raji, COS, QT35, BL3. Similar results were obtained in cell adhesion assay using MN-Fc protein dropped on
25 bacterial Petri dishes. These assays revealed that KATO III human stomach adenocarcinoma cell line is lacking an ability to interact with MN-Fc protein. This finding allowed us to use KATO III cells for expression cloning and screening of the cDNA coding for MN-binding protein.

The cDNA expression library in pBK-CMV vector was prepared from
30 dense HeLa cells and used for transfection of KATO III cells. For the first round of screening, KATO III cells were transfected by electroporation. After two days of incubation, the ligand-expressing cells were allowed to bind to MN-Fc protein, then to

Protein A conjugated with biotin and finally selected by pulling down with streptavidin-coated magnetic beads. Plasmid DNA was extracted from the selected cells and transformed to E. coli. Individual E. coli colonies were picked and pools of 8-10 clones were prepared. Plasmid DNA from the pools was isolated and used in the second
5 round of screening.

In the second round of screening, KATO III cells were transfected by DEAE dextran method. To identify the pool containing the cDNA for MN-binding protein, an ELISA method based on the binding of MN-Fc to the transfected cells, and detection using peroxidase labelled Protein A were used. Pools are selected by ability
10 to bind MN-Fc.

In the third round of screening, plasmid DNAs isolated from individual bacterial colonies of selected pools are transfected to KATO III cells. The transfected cells are subjected to binding with MN-Fc and detection with Protein A as before. Such exemplary screening is expected to identify a clone containing the cDNA which codes
15 for the putative MN protein partner. That clone would then be sequenced and the expression product confirmed as binding to MN protein by cell adhesion assay. (Far-Western blotting, co-precipitation etc.) Hybridomas producing Mabs to the expression product would then be prepared which would allow the analysis of the biological characteristics of the protein partner of MN.

20

Preparation of MN-Specific Antibodies

The term "antibodies" is defined herein to include not only whole antibodies but also biologically active fragments of antibodies, preferably fragments containing the antigen binding regions. Further included in the definition of antibodies
25 are bispecific antibodies that are specific for MN protein and to another tissue-specific antigen.

Zavada et al., WO 93/18152 and WO 95/34650 describe in detail methods to produce MN-specific antibodies, and detail steps of preparing representative MN-specific antibodies as the M75, MN7, MN9, and MN12 monoclonal
30 antibodies. Preferred MN antigen epitopes comprise: aa 62-67 (SEQ ID NO: 10); aa 61-66, aa 79-84, aa 85-90 and aa 91-96 (SEQ ID NO: 98); aa 62-65, aa 80-83, aa 86-89 and aa 92-95 (SEQ ID NO: 99); aa 62-66, aa 80-84, aa 86-90 and aa 92-96 (SEQ ID

NO: 100); aa 63-68 (SEQ ID NO: 101); aa 62-68 (SEQ ID NO: 102); aa 82-87 and aa 88-93 (SEQ ID NO: 103); aa 55-60 (SEQ ID NO: 11); aa 127-147 (SEQ ID NO: 12); aa 36-51 (SEQ ID NO: 13); aa 68-91 (SEQ ID NO: 14); aa 279-291 (SEQ ID NO: 15); and aa 435-450 (SEQ ID NO: 16). Example 2 provides further description concerning preferred MN antigen epitopes.

Bispecific Antibodies. Bispecific antibodies can be produced by chemically coupling two antibodies of the desired specificity. Bispecific MAbs can preferably be developed by somatic hybridization of 2 hybridomas. Bispecific MAbs for targeting MN protein and another antigen can be produced by fusing a hybridoma that produces MN-specific MAbs with a hybridoma producing MAbs specific to another antigen. For example, a cell (a quadroma), formed by fusion of a hybridoma producing a MN-specific MAb and a hybridoma producing an anti-cytotoxic cell antibody, will produce hybrid antibody having specificity of the parent antibodies. [See, e.g., Immunol. Rev. (1979); Cold Spring Harbor Symposium Quant. Biol., 41: 793 (1977); van Dijk et al., Int. J. Cancer, 43: 344-349 (1989).] Thus, a hybridoma producing a MN-specific MAb can be fused with a hybridoma producing, for example, an anti-T3 antibody to yield a cell line which produces a MN/T3 bispecific antibody which can target cytotoxic T cells to MN-expressing tumor cells.

It may be preferred for therapeutic and/or imaging uses that the antibodies be biologically active antibody fragments, preferably genetically engineered fragments, more preferably genetically engineered fragments from the V_H and/or V_L regions, and still more preferably comprising the hypervariable regions thereof. However, for some therapeutic uses bispecific antibodies targeting MN protein and cytotoxic cells would be preferred.

25

Epitopes

The affinity of a MAb to peptides containing an epitope depends on the context, e.g. on whether the peptide is a short sequence (4-6 aa), or whether such a short peptide is flanked by longer aa sequences on one or both sides, or whether in testing for an epitope, the peptides are in solution or immobilized on a surface. Therefore, it would be expected by ones of skill in the art that the representative

30

epitopes described herein for the MN-specific MABs would vary in the context of the use of those MABs.

The term "corresponding to an epitope of an MN protein/polypeptide" will be understood to include the practical possibility that, in some instances, amino acid sequence variations of a naturally occurring protein or polypeptide may be antigenic and confer protective immunity against neoplastic disease and/or anti-tumorigenic effects. Possible sequence variations include, without limitation, amino acid substitutions, extensions, deletions, truncations, interpolations and combinations thereof. Such variations fall within the contemplated scope of the invention provided the protein or polypeptide containing them is immunogenic and antibodies elicited by such a polypeptide or protein cross-react with naturally occurring MN proteins and polypeptides to a sufficient extent to provide protective immunity and/or anti-tumorigenic activity when administered as a vaccine.

Epitope for M75 MAb

The M75 epitope is considered to be present in at least two copies within the 6X tandem repeat of 6 amino acids [aa 61-96 (SEQ ID NO: 97)] in the proteglycan domain of the MN protein. Exemplary peptides representing that epitope depending on the context may include the following peptides from that tandem repeat: EEDLPS (SEQ ID NO: 10; aa 62-67); GEEDLP (SEQ ID NO: 98; aa 61-66; aa 79-84; aa 85-90; aa 91-96); EEDL (SEQ ID NO: 99; aa 62-65; aa 80-83; aa 86-89; aa 92-95); EEDLP (SEQ ID NO: 100; aa 62-66; aa 80-84; aa 86-90; aa 92-96); EDLPSE (SEQ ID NO: 101; aa 63-68); EEDLPSE (SEQ ID NO: 102; aa 62-68); and DLPGEE (SEQ ID NO: 103; aa 82-87, aa 88-93).

Three synthetic peptides from the deduced aa sequence for the EC domain of the MN protein shown in Figure 1 were prepared. Those synthetic peptides are represented by aa 51-72 (SEQ ID NO: 104), aa 61-85 (SEQ ID NO: 105) and aa 75-98 (SEQ ID NO: 106). Each of those synthetic peptides contains the motif EEDLP (SEQ ID NO: 100) and were shown to be reactive with the M75 MAb.

Other Epitopes

Mab MN9. Monoclonal antibody MN9 (Mab MN9) reacts to the same epitope as Mab M75, as described above. As Mab M75, Mab MN9 recognizes both the GST-MN fusion protein and native MN protein equally well.

5 Mabs corresponding to Mab MN9 can be prepared reproducibly by screening a series of mabs prepared against an MN protein/polypeptide, such as, the GST-MN fusion protein, against the peptides representing the epitope for Mabs M75 and MN9. Alternatively, the Novatope system [Novagen] or competition with the deposited Mab M75 could be used to select mabs comparable to Mabs M75 and MN9.

10 Mab MN12. Monoclonal antibody MN12 (Mab MN12) is produced by the mouse lymphocytic hybridoma MN 12.2.2 which was deposited under ATCC HB 11647. Antibodies corresponding to Mab MN12 can also be made, analogously to the method outlined above for Mab MN9, by screening a series of antibodies prepared against an MN protein/polypeptide, against the peptide representing the epitope for
15 Mab MN12. That peptide is aa 55 - aa 60 of Figure 1 [SEQ ID NO: 11]. The Novatope system could also be used to find antibodies specific for said epitope.

Mab MN7. Monoclonal antibody MN7 (Mab MN7) was selected from mabs prepared against nonglycosylated GST-MN as described above. It recognizes the epitope represented by the amino acid sequence from aa 127 to aa 147 [SEQ ID NO:
20 12] of the Figure 1 MN protein. Analogously to methods described above for Mabs MN9 and MN12, mabs corresponding to Mab MN7 can be prepared by selecting mabs prepared against an MN protein/polypeptide that are reactive with the peptide having SEQ ID NO: 12, or by the stated alternative means.

25 MN-Specific Intrabodies – Targeted Tumor Killing Via Intracellular
 Expression of MN-Specific Antibodies to
 Block Transport of MN Protein to Cell Surface

 The gene encoding antibodies can be manipulated so that the antigen-
30 binding domain can be expressed intracellularly. Such “intrabodies” that are targeted to the lumen of the endoplasmic reticulum provide a simple and effective mechanism for inhibiting the transport of plasma membrane proteins to the cell surface. [Marasco, W.A., “Review – Intrabodies: turning the humoral immune system outside in or

intracellular immunization," Gene Therapy, 4: 11-15 (1997); Chen et al., "Intracellular antibodies as a new class of therapeutic molecules for gene therapy," Hum. Gene Ther., 5(5): 595-601 (1994); Mhashilkar et al., EMBO J., 14: 1542-1551 (1995); Mhashilkar et al., J. Virol., 71: 6486-6494 (1997); Marasco (Ed.), Intrabodies: Basic Research and Clinical Gene Therapy Applications, (Springer Life Sciences 1998; ISBN 3-540-64151-3) (summarizes preclinical studies from laboratories worldwide that have used intrabodies); Zanetti and Capra (Eds.), "Intrabodies: From Antibody Genes to Intracellular Communication," The Antibodies: Volume 4, [Harwood Academic Publishers; ISBN 90-5702-559-0 (Dec. 1997)]; Jones and Marasco, Advanced Drug Delivery Reviews, 31 (1-2): 153-170 (1998); Pumphrey and Marasco, Biodrugs, 9(3): 179-185 (1998); Dachs et al., Oncology Res., 9(6-7): 313-325 (1997); Rondon and Marasco, Ann. Rev. Microbiol., 51: 257-283 (1997)]; Marasco, W.A., Immunotechnology, 1(1): 1-19 (1995); and Richardson and Marasco, Trends in Biotechnology, 13(8): 306-310 (1995).]

15 MN-specific intrabodies may prevent the maturation and transport of MN protein to the cell surface and thereby prevent the MN protein from functioning in an oncogenic process. Antibodies directed to MN's EC, TM or IC domains may be useful in this regard. MN protein is considered to mediate signal transduction by transferring signals from the EC domain to the IC tail and then by associating with other intracellular
20 proteins within the cell's interior. MN-specific intrabodies could disrupt that association and perturb that MN function.

Inactivating the function of the MN protein could result in reversion of tumor cells to a non-transformed phenotype. [Marasco et al. (1997), supra.] Antisense expression of MN cDNA in cervical carcinoma cells, as demonstrated herein, has
25 shown that loss of MN protein has led to growth suppression of the transfected cells. It is similarly expected that inhibition of MN protein transport to the cell surface would have similar effects. Cloning and intracellular expression of the M75 MAb's variable region is to be studied to confirm that expectation.

Preferably, the intracellularly produced MN-specific antibodies are single-
30 chain antibodies, specifically single-chain variable region fragments or sFv, in which the heavy- and light-chain variable domains are synthesized as a single polypeptide and are separated by a flexible linker peptide, preferably (Gly₄-Ser)₃ [SEQ ID NO: 116].

MN-specific intracellularly produced antibodies can be used therapeutically to treat preneoplastic/neoplastic disease by transfecting preneoplastic/neoplastic cells that are abnormally expressing MN protein with a vector comprising a nucleic acid encoding MN-specific antibody variable region fragments, 5 operatively linked to an expression control sequence. Preferably said expression control sequence would comprise the MN gene promoter.

Antibody-Mediated Gene Transfer Using MN-Specific
Antibodies or Peptides for Targeting MN-Expressing Tumor Cells

10 An MN-specific antibody or peptide covalently linked to polylysine, a polycation able to compact DNA and neutralize its negative charges, would be expected to deliver efficiently biologically active DNA into an MN-expressing tumor cell. If the packed DNA contains the HSVtk gene under control of the MN promoter, the system would have double specificity for recognition and expression only in MN- 15 expressing tumor cells. The packed DNA could also code for cytokines to induce CTL activity, or for other biologically active molecules. The M75 MAb (or, for example, as a single chain antibody, or as its variable region) is exemplary of such a MN-specific antibody.

The following examples are for purposes of illustration only and are not 20 meant to limit the invention in any way.

Example1

Transient Transformation of
Mammalian Cells by MN Protein

25 This example (1) examines the biological consequences of transfecting human or mouse cells with MN-cDNA inserted into expression vectors, mainly from the viewpoint of the involvement of MN protein in oncogenesis; (2) determines if MN protein exerts carbonic anhydrase activity, and whether such activity is relevant for 30 morphologic transformation of cells; and (3) tests whether MN protein is a cell adhesion molecule (CAM).

Synopsis

Methods: MN-cDNA was inserted into 3 expression vectors and was used for transfecting human or mouse cells. MN protein was detected by Western blotting, radioimmunoassay or immunoperoxidase staining; in all tests the MN-specific
5 monoclonal antibody M75 (MAb M75) was used. Carbonic anhydrase activity was determined by the acidification velocity of carbonate buffer in CO₂ atmosphere.

Results: (1) Cells (human CGL-1 and mouse NIH3T3 cells) transfected with MN-cDNA showed morphologic transformation, but reverted to normal phenotype after 4-5 weeks. (2) This reversion was not due to the loss, silencing or mutation of the
10 MN insert. (3) MN protein has the enzyme activity of a carbonic anhydrase, which can be inhibited with acetazolamide; however, the inhibition of the carbonic anhydrase enzyme activity did not affect transformation. (4) MN protein is an adhesion protein, involved in cell-to-cell contacts.

15 Background

This example concerns transformation of mammalian cells by MN-cDNA inserted into expression vectors derived from retroviruses. Such vectors are suitable for efficient and stable integration into cellular DNA and for continuous expression of MN protein. Cells transfected with these constructs showed morphologic transformation,
20 but after some time, they reverted to normal phenotype.

Sulfonamides, including acetazolamide, are very potent inhibitors of known carbonic anhydrases [Maren and Ellison, Mol. Pharmacol., 3: 503-508 (1967)]. Acetazolamide was tested to determine if it inhibited also the MN-carbonic anhydrase, and if so, whether inhibition of the enzyme affected cell transformation.

25 There are reasons to believe that MN protein could be involved in direct cell-to-cell interactions: A) previous observations indicated a functional resemblance of MN protein to surface glycoproteins of enveloped viruses, which mediate virus adsorption to cell surface receptors, and MN participated in the formation of phenotypically mixed virions of vesicular stomatitis virus. B) Inducibility of MN protein
30 expression by growing HeLa cells in densely packed monolayers suggests that it may be involved in direct interactions between cells. C) Finally, there is a structural similarity between the MN protein and receptor tyrosine phosphatase β , which also contains

proteoglycan and carbonic anhydrase domains; those domains mediate direct contacts between cells of the developing nervous system [Peles et al., Cell, 82: 251-260 (1995)]. Therefore, MN protein was tested to see if it bound to cell surface receptors; the result was clearly positive that it does.

5

Materials and Methods

Cell Lines

Cells used in this example were: CGL1 and CGL3 - respectively non-tumorigenic and tumorigenic HeLa x fibroblast hybrids [Stanbridge et al., Somat. Cell Genet., 7: 699-712 (1981)], mouse cell line NIH3T3, HeLa cells and monkey Vero cells. The NIH3T3 cells were seeded at very low density to obtain colonies started from single cells. The most normal appearing colony, designated subclone 2, was picked for use in the experiments reported in this example.

15

Expression Vectors

Full-length MN cDNA was acquired from a pBluescript subclone [Pastorek et al., Oncogene, 9: 2877-2888 (1994)]. To remove 5' and 3' noncoding sequences, that might reduce subsequent gene expression, a polymerase chain reaction (PCR) was performed. The 5' primer
TAGACAGATCTACGATGGCTCCCCTGTGCCCCAG [SEQ ID NO: 88] encompasses a translation start site and BglII cloning site, and the 3' primer
ATTCCTCTAGACAGTTACCGGCTCCCCCTCAGAT [SEQ ID NO: 89] encompasses a stop codon and XbaI cloning site. Full-length MN-cDNA as a template and Pfu DNA Polymerase [Stratagene; LaJolla, CA (USA)] were used in the reaction.

25

The PCR product was sequenced and found to be identical with the template; it carried no mutations. The PCR product harbouring solely the MN coding sequence was inserted into three vectors: 1. pMAMneo [Clontech; Palo Alto, CA (USA)] plasmid allowing dexamethasone-inducible expression driven by the MMTV-Long Terminal Repeat (LTR) promoter and containing a neo gene for selection of
transformants in media supplemented with Geneticin (G418) antibiotics. 2. Retroviral expression vector pGD [Daley et al., Science, 247: 824-829 (1990); kindly provided by
Prof. David Baltimore, New York-Cambridge)] containing MLV-LTR promoter and neo

30

gene for G418 antibiotics selection. 3. Vaccinia virus expression vector pSC11 [Chakrabarti et al., Mol. Cell. Biol., 5: 3403-3409 (1985)]. Transfection was performed via a calcium-phosphate precipitate according to Sambrook et al. (eds.), Molecular cloning. A laboratory manual, 2nd ed., Cold Spring Harbor Laboratory Press (1989).

5 Vaccinia virus strain Praha clone 13 was used as parental virus [Kutinova et al., Vaccine, 13: 487-493 (1995)]. Vaccinia virus recombinant was prepared by a standard procedure [Perkus et al., Virology, 152: 285-297 (1986)]. Recombinant viruses were selected and plaque purified twice in rat thymidine-kinase-less RAT2 cells [Topp, W. C., Virology, 113: 408-411 (1981)] in the presence of 5'-bromodeoxyuridine
10 (100 μ g/ml). Blue plaques were identified by overlaying with agar containing 5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside (X-Gal) (200 μ g/ml).

CA Assay

Carbonic anhydrase activity was measured by a micro-method [Brion et al., Anal. Biochem., 175: 289-297 (1988)]. In principle, velocity of the reaction $\text{CO}_2 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{CO}_3$ is measured by the time required for acidification of carbonate buffer, detected with phenol red as a pH indicator. This reaction proceeds even in absence of the enzyme, with t_0 = control time (this was set to 60 seconds). Carbonic anhydrase reduces the time of acidification to t ; one unit of the enzyme activity reduces the time
20 to one half of control time: $t/t_0 = 1/2$.

For the experiment, MN protein was immunoprecipitated with Mab M75 from RIPA buffer (1% Triton X-100, 0.1% deoxycholate, 1mM phenylmethylsulfonyl-fluoride and 200 trypsin-inhibiting units/ml of Trasylol in PBS, pH 7.2) extract of Vero cells infected with vaccinia-MN construct, after the cells developed cytopathic effect, or
25 with "empty" vaccinia as a control. The MN + antibody complex was subsequently adsorbed to protein A - Staphylococcus aureus cells [Kessler, S. W., J. Immunol., 115: 1617-1624 (1975)] and rinsed 2x with PBS and 2x with 1 mM carbonate buffer, pH 8.0. The precipitate was resuspended in the same buffer and added to the reaction mixture. Acetazolamide (Sigma) was tested for inhibition of carbonic anhydrase [Maren and
30 Ellison, supra]. In extracts of infected cells used for immunoprecipitation, the concentration of total proteins was determined by the Lowry method [Lowry et al., J.

Biol. Chem., 193: 265-275 (1951)] and that of MN protein by a competition radioimmunoassay as described in Zavada et al., Int. J. Cancer, 54: 268-274 (1993).

Western Blots

5 Western blotting and development of the blots using ¹²⁵I-labelled M75 and autoradiography was performed as before [Pastorekova et al., Virology, 187: 620-626 (1992); and Zavada (1993), supra].

Adhesion Assay

10 For the adhesion assay [Hoffman S., "Assays of cell adhesion," IN: Cell-cell Interactions, (Stevenson et al. eds.) pp. 1-30 (IRL Press at Oxford University Press; Oxford, N.Y., Tokyo; 1992)], 25 μ l aliquots MN protein (affinity purified pGEX-3X MN) [Zavada et al. (1993), supra] or of control proteins were spotted on 5 cm-diameter bacteriological Petri dishes and allowed to bind for 2 hours at room temperature. This
15 yielded circular protein-coated areas of 4-5 mm diameter. MN protein was diluted to 10 μ g/ml in 50 mM carbonate buffer, pH 9.2. Patches of adsorbed control proteins were prepared similarly. Those included collagens type I and IV, fibronectin, laminin and gelatin (Sigma products), diluted and adsorbed according to the manufacturer's recommendations; FCS and BSA were also included. After aspiration of the drops, the
20 dishes were rinsed 2x with PBS and saturated for 1 hour with DMEM supplied with 5% FCS. The plates were seeded with 5×10^5 cells in 5 ml of DMEM + 5% FCS and incubated overnight at 37°C. The plates were rinsed with PBS, and the attached cells were fixed with formaldehyde, post-fixed with methanol and Giemsa stained.

25 Results

1. Transformation and reversion of CGL1 cells transfected with MN-cDNA

Since the expression of MN protein correlated with the tumorigenicity of HeLa x fibroblast hybrids [Zavada et al. (1993), supra], the non-tumorigenic hybrid
30 CGL1 cells were first tested. Those cells, transfected with the pMAM.MN construct, after selection with Geneticin, formed colonies with varying degrees of transformation; some of them appeared normal. While normal CGL1 cells are contact inhibited, growing in a parallel orientation, the transformed cells formed very dense colonies,

showing the loss of contact inhibition. Such colonies grew more slowly than the original CGL 1.

After subcloning, the cells isolated from transformed colonies segregated revertants. The reversion was a gradual, step-wise process; there were colonies with
5 different degrees of reversion. After 2 passages, all the cell population became a morphologically indistinguishable from normal CGL1. This was due to the reversion of some cells and to the selective advantage of the revertants, which grew faster than the transformed cells. Despite repeated attempts, not even one single stably transformed cell clone was obtained. No transformed colonies were found in CGL1 cells
10 transfected with an "empty" pMAM control plasmid. Growth of the CGL1 + pMAM.MN revertants in media supplied with 5 μ g/ml of dexamethasone for 7 days enhanced the production of MN protein, but the morphology of the cells did not return to transformed.

15 2. Rescue of transforming MN from the revertants

The reversion of MN-transformed cells to normal phenotype could have at least 4 causes: A) loss of the MN insert; B) silencing of the MN insert, e.g., by methylation; C) mutation of the MN insert; D) activation of a suppressor gene, coding for a product which neutralizes transforming activity of MN protein; E) loss of a MN-
20 binding protein. To decide among those alternatives, the following experiment was designed.

MN-cDNA was inserted into pGD, a vector derived from mouse leukemia virus - MLV. A defective virus was thereby engineered, which contained the MN gene and the selective marker neo instead of genes coding for viral structural proteins. With
25 this construct, mouse NIH3T3 cells were transfected. In media supplied with Geneticin, the cells formed colonies with phenotypes ranging from strongly transformed to apparently normal. All of the transformed colonies and about 50% of the normal colonies expressed MN protein. Contrasting with normal NIH3T3 cells, the transformants were also able to form colonies in soft agar, reflective of the loss of
30 anchorage dependence, characteristic of cell transformation. Upon passaging, the cells isolated from transformed colonies reverted to normal morphology, and at the same time, they lost the capacity to form colonies in soft agar, while still expressing the MN

protein. This permanent presence of MN protein in revertants ruled out alternatives A) and B) supra, that is, loss or silencing of the MN gene as a cause of reversion.

To decide among the other 3 alternatives, the revertants were superinfected with live, replication competent MLV. This virus grows in NIH3T3 cells without any morphologic manifestations, and it works as a "helper" for the pGD.MN construct. Virus progeny from MLV-infected revertants represents an artificial virus complex [pGD.MN + MLV]. This consists of 2 types of virions: of standard type MLV particles and virions containing the pGD.MN genome, enveloped in structural proteins provided by the "helper" virus. This virus complex was infectious for fresh NIH3T3 cells; it again induced in them morphologic transformation and the capacity to form agar colonies.

Contrasting with NIH3T3 transfected with pGD.MN, all the colonies of cells infected with [pGD.MN + MLV] complex, which grew in the presence of Geneticin, were uniformly transformed and contained MN proteins. The transformants once more reverted to normal phenotype although they kept producing infectious [pGD.MN + MLV] complex, which induced transformation in fresh NIH3T3 cells. This cycle of infection-transformation-reversion was repeated 3 times with the same result. This ruled out alternative C) – mutation of MN-cDNA as a cause of reversion.

Normal NIH3T3 cells formed a contact inhibited monolayer of flat cells, which did not stain with Mab M75 and immunoperoxidase. Cells infected with [pGD.MN + MLV] complex were clearly transformed: they grew in a chaotic pattern and showed loss of contact inhibition. Some of the cells showed signs of apoptosis. Two passages later, the cell population totally reverted to original phenotype as a result of frequent emergence of revertants and of their selective advantages (faster growth and a higher efficiency of plating). In fact, the revertants appeared to grow to a somewhat lower saturation density than the original NIH3T3 cells, showing a higher degree of contact inhibition.

The control NIH3T3 cells did not contain any MN protein (Western blot); while both transformed cells and revertants contained the same amount and the same proportion of 54 and 58 kDa bands of MN protein. In a non-reducing gel, MN protein was present in the form of oligomers of 153 kDa. Consistently, by competition RIA,

approximately 40 ng MN/mg total protein was found in both of the transformed cells and revertants.

3. Carbonic anhydrase activity and its inhibition

5 Since the carbonic anhydrase domain represents a considerable part of the MN protein (see Figure 8), tests were performed to determine whether it is indeed enzymatically active. Vero cells infected with the vaccinia.MN construct, which contained more of the MN protein than other cells used in the present experiments, served as a source of MN protein. The cells were extracted with RIPA buffer, and MN
10 protein was concentrated and partially purified by precipitation with MAb M75 and SAC. The immunoprecipitate was tested for CA activity. 78 μ l of precipitate contained 1 unit of the enzyme. From the extract, the concentration of total proteins and of MN protein was determined; 1 unit of enzyme corresponded to 145 ng of MN protein or to 0.83 mg of total protein. The immunoprecipitate from Vero cells infected with control
15 virus had no enzyme activity. Activity of MN carbonic anhydrase was inhibited by acetazolamide; 1.53×10^{-8} M concentration of the drug reduced enzyme activity to 50%.

 Preliminary tests showed that confluent cultures of HeLa or of NIH3T3 cells tolerated 10^{-5} - 10^{-3} M concentration of acetazolamide for 3 days without any signs
20 of toxicity and without any effect on cell morphology. In sparse cultures, 10^{-5} M acetazolamide did not inhibit cell growth, but 10^{-4} M already caused a partial inhibition. Thus, 10^{-5} M acetazolamide was added to NIH3T3 cells freshly transformed with the [pGD.MN + MLV] complex. After 4 days of incubation, the colonies were fixed and stained. No difference was seen between cells growing in the presence or
25 absence of acetazolamide; both were indistinguishable from correctly transformed NIH3T3 cells. Thus, the enzymatic activity of carbonic anhydrase is not relevant for the transforming activity of MN protein.

4. Cell adhesion assay

30 To determine whether or not MN protein is a cell adhesion molecule (CAM), adhesion assays were performed in plastic bacteriological Petri dishes (not treated for use with tissue culture). Cells do not adhere to the surfaces of such dishes,

unless the dishes are coated with a binding protein. NIH3T3 cells adhered, spread and grew on patches of adsorbed MN protein. Only very few cells attached outside the areas coated with MN protein.

Other variants of the experiment demonstrated that NIH3T3 cells adhered and spread on patches of adsorbed collagen I and IV, fibronectin and laminin. NIH3T3 cells did not attach to dots of adsorbed gelatin, FCS or BSA.

CGL1, HeLa and Vero cells also adhered to MN protein, but 3 leukemia cell lines showed no adherence. CGL3 cells, strongly expressing MN protein adhered less efficiently to MN protein dots than did CGL1. The presence of 10^{-4} M acetazolamide in the media did not affect the cell adhesion.

To confirm the specificity of adhesion, MN protein was absorbed with SAC loaded with MAb M75 (directed to MN) or MAb M67, directed to an unrelated antigen (Pastorekova et al., *supra*), before it was applied to the surface of the Petri dishes. Absorption with the SAC-M75 complex totally abrogated the cell binding activity, whereas absorption with SAC-M67 was without any effect.

Additional Cell Adhesion Results

A shortened MN, missing TM and IC segments, is shed into the medium by 5ET1 cells (a HeLa X fibroblast hybrid, analogous to CGL3 cells that express MN protein abundantly) or by Vero cells infected with VV carrying MN-cDNA with deleted TM and IC sequences. The shed MN protein was purified from the media, and tested in cell adhesion assays. The cells adhered, spread and grew only on the patches covered with adsorbed complete MN protein, but not on the dots of MN lacking TM and IC regions. Analogous results have been described also for some other adhesion molecules. A variety of cells (NIH3T3, CGL1, CGL3, HeLa, XC) attached to MN protein dots suggesting that the MN receptor(s) is common on the surface of vertebrate cells.

Tests were also performed with extracellular matrix proteins or control proteins dotted on nitrocellulose. The dot-blots were treated with MN protein solution. Bound MN protein was detected with MAb M75. MN protein adsorbed to the dots of collagen I and IV, but not to fibronectin, laminin, gelatine or BSA.

Prospects for therapy. There are many new principles of cancer therapy employing oncoproteins or molecules that interact with them as targets [Mendelsohn and Lippman, "Principles of molecular cell biology of cancer: growth factors," In: DeVita et al., eds., Cancer: principles and practice of oncology, pp. 114-133 4th ed., Philadelphia: Lippincott (1993); DeVita et al., eds., Biologic therapy of cancer, 2nd ed., Philadelphia: Lippincott (1995)]. The MN protein and at least some of its ligands (or receptors) appear to be particularly suitable for such purposes.

Example 2

Identification of MN's Binding Site

MN protein is a tumor-associated cell adhesion molecule (CAM). To identify its binding site, a series of overlapping oligopeptides, spanning the N-terminal domain of the MN protein were synthesized. The N-terminal domain is homologous to that of proteoglycans and contains a tandem repeat of six amino acids.

The series of oligopeptides were tested by the cell adhesion assay procedure essentially as described above in Example 1. The synthetic oligopeptides were immobilized on hydrophobic plastic surfaces to see if they would mediate the attachment, spreading and growth of cells. Also investigated were whether the oligopeptides or antibodies inhibited attachment of cells (NIH3T3, HeLa and CGL1) to purified MN protein coated onto such plastic surfaces. The MN protein was affinity purified on agarose covalently linked to sulfonamide, as the MN protein encompasses a CA domain.

Several of the oligopeptides were found to be biologically active: (i) when immobilized onto the plastic, they mediate attachment of cells (NIH3T3, HeLa and to CGL1); (ii) when added to the media, they compete for attachment to cells with the immobilized MN protein; (iii) these oligopeptides, present in the media do not inhibit attachment of cells to TC plastic, but they prevent cell-cell adhesion and formation of intercellular contacts; (iv) treatment of immobilized MN protein and of active peptides with MAb M75 abrogates their affinity for the cells; and (v) the binding site of MN was determined to be closely related or identical to the epitope for MAb M75, at least two copies of which are located in the 6-fold tandem repeat of 6 amino acids [aa 61-96 (SEQ ID NO: 97)] in the proteoglycan-like domain of MN protein.

It was concluded that ectopically expressed MN protein most likely participates in oncogenesis by intervention into normal cell-cell contacts. MN's binding site represents a potential target for which therapeutic agents can be designed.

5

Materials and Methods

Affinity chromatography of MN/CA IX. MN/CA IX was purified by a single cycle of adsorption - elution on sulfonamide-agarose, as described for other CAs [Falkbring et al., FEBS Letters, 24: 229 (1972)]. We used columns of p-aminoethylbenzenesulfonamide-agarose (Sigma). Columns with adsorbed MN/CA IX were extensively washed with PBS (NaCl 8.0 g/l, KCl 0.2 g/l, KH₂PO₄ 0.2 g/l, Na₂HPO₄ 1.15 g/l, pH=7.2) and eluted with 0.1 mM acetazolamide (Sigma). All steps of purification were carried out at 0 - 5 °C, pH 7.2, at physiological concentration of salts. Complete MN/CA IX+ was extracted with 1% Triton X-100 in PBS from Vero cells infected with vaccinia virus containing an insert of complete coding region of MN/CA IX as described in Zavada et al., Int. J. Oncol., 10: 857 (1997). Before chromatography, the extract was diluted 1:6 with PBS and centrifuged for 1 h at 1500 xg. Truncated MN/CA IX Δ TM Δ IC was produced from an analogous construct except that the 3' downstream primer for PCR was: 5' CGT CTA GAA GGA ATT CAG CTA GAC TGG CTC AGC A 3' [SEQ ID NO: 117]. MN/CA IX Δ was shed into the medium, from which it was affinity purified after centrifugation as above. All steps of purification were monitored by dot-blot.

Cells and media. The following cell lines were used: HeLa, CGL1 = non-tumorigenic hybrid HeLa x fibroblast, CGL3 = tumorigenic segregant from this hybrid, NIH3T3 cells = mouse fibroblasts. The origin of the cells and growth media are described in Zavada et al., Int. J. Cancer, 54: 268 (1993) and Zavada et al., Int. J. Oncol., 10: 857 (1997). In addition, we used also HT29, a cell line derived from colorectal carcinoma (ATCC No. HBT-38).

Cell adhesion assay. The conditions of the assay are basically as described in Example 1. Briefly, 1 μ g/ml of purified MN/CA IX in 50 mM mono/bicarbonate buffer, pH 9.2, was adsorbed in 30 μ l drops on the bottom of bacteriological 5 cm Petri dishes for 1.5 hr. Then the drops were removed by aspiration and the dishes were 3x rinsed with PBS and blocked with 50% FCS in culture medium for 30 min. There were two

variants of the test. In the first one, the whole bottom of the Petri dish was blocked with 50% FCS, and the dishes were seeded with 5 ml of cell suspension (10^5 cells/ml). After overnight incubation, the cultures were rinsed with PBS, fixed and stained. In the other variant, only the area of adsorbed MN/CA IX was blocked and on top of MN/CA IX dots were added 30 μ l drops of cell suspension in growth medium, containing added oligopeptides (or control without peptides). After incubation, rinsing and fixation, the cultures were stained with 0.5% Trypan blue in 50 mM Tris buffer pH 8.5 for 1 h, rinsed with water and dried. Stained areas of attached cells were extracted with 10% acetic acid, the extracts transferred to 96-well plates and absorbance was measured at 630 nm on microplate reader.

ELISA. Purified GST-MN [Zavada et al. (1993), *supra*] at concentration 10 ng/ml in carbonate buffer pH 9.2 was adsorbed for 3 h in Maxisorb strips (NUNC). After washing and blocking (1 h) with 0.05% Tween 20 in PBS, 50 μ l/well of the antibody + antigen mixtures were added. Final dilution of MAb 75 ascites fluid was 10^{-6} ; concentration of the peptides varied according to their affinity for M75 so as to allow determination of 50% end-point. These mixtures were adsorbed for 1.5 h, followed by washing with Tween-PBS. Bound antibody was detected by antimouse IgG conjugate with peroxidase (SwAM-Px, SEVAC, Prague), diluted 1:1000. In the color reaction OPD (o-phenylenediamine dihydrochloride, Sigma) 1mg/ml in 0.1 M citrate buffer pH 5.0 was used. To this H_2O_2 was added to final concentration 0.03%. This system is balanced so as to allow assay for antigen competing for M75 as well as for peptides binding to the epitope of immobilized GST-MN.

Peptides. The peptides used in this study were prepared by the solid phase method [Merrifield et al., *IN: Gutte, B. (ed.), Peptides: Synthesis, Structures and Applications*, pp. 93-169 (San Diego; Academic Press; 1995)] using the Boc/Bzl strategy. The peptide acids were prepared on PAM-resin and peptide amides on MeBHA resin. Deprotection and splitting from the resin was done by liquid hydrogen fluoride. The peptides were purified by C18 RP HPLC and characterized by amino acid analysis and FAB MS spectroscopy.

Western blots. MN/CA IX antigens from PAGE gels were transferred to PVDF membranes (Immobilon P, Millipore) and developed with M75, followed by

SwAM-Px (see above) and diaminobenzidine (Sigma) with H₂O₂. For dot-blots we used nitrocellulose membranes.

Phage display. Ph.D.-7 Phage Display Peptide Library kit was used for screening as recommended by manufacturer (New England Biolabs). 96-well plate was coated with peptide SEQ ID NO: 106. Biopanning was carried out by incubating 2x10¹¹ phage with target coated plate for 1 h. Unbound phages were washed away with TBST (50mM Tris-HCl pH 7.5, 150 mM NaCl, 0.1% Tween-20) and specifically bound phages were eluted with M75 antibody (2μg in 100 μl of TBS/well). Eluted phage was amplified and used for additional binding and amplification cycles to enrich the pool in favour of binding sequence. After 5 rounds, individual clones were picked, amplified and sequenced using T7 sequencing kit (Pharmacia).

Results

Affinity chromatography of MN/CA IX protein. For purification of MN/CA IX protein we decided to use affinity chromatography on sulfonamide-agarose column, described previously for other CAs [Falkbring et al., *supra*]. The advantages of this method are simplicity and the fact that the whole procedure is carried out under non-denaturing conditions. Vaccinia virus vector with an insert of the complete MN/CA9 cDNA, or with truncated cDNA (lacking transmembrane and intracellular domains) was employed as a source of MN/CA IX protein.

A single cycle of adsorption - elution yielded relatively pure proteins: MN/CA IX+ gave 2 bands of 54 and 58 kDa, MN/CA IXΔ of 54.5 and 56 kDa. These proteins strongly reacted with MAb M75 on Western blots. In extracts from HeLa, CGL3 and HT29 the blot revealed 2 bands of the same size as MN/CA IX+ purified from vaccinia virus construct.

Adhesion of cells to MN/CA IX protein. MN/CA IX immobilized on hydrophobic plastic enabled attachment, spreading and growth of cells. Extremely low concentrations of MN/CA IX corresponding to 1 μg/ml of purified protein in adsorption buffer were sufficient to cause this effect; other cell adhesion molecules are used in 10 - 50x higher concentrations. Only complete MN/CA IX protein was active in cell adhesion test, truncated MN/CA IX did not support cell adhesion at all or it showed only a low adhesion activity and in some instances it even acted as a cell "repellent".

Treatment of the dots of immobilized MN/CA IX with MAb M75 abrogated its capacity to attach the cells, but the control MAb M16, irrelevant for MN/CA IX had no effect. Blocking of cell attachment by M75 shows that the epitope is identical to or overlapping with the binding site of MN/CA IX for cell receptors.

- 5 Identification of the epitope recognized by Mab M75. Preliminary mapping of M75 epitope employing partial sequences of extracellular parts of MN/CA9 cDNA expressed from bacterial vectors and tested on Western blots located it in PG region. For exact mapping, our strategy was to synthesize partially overlapping oligopeptides of 15-25 aa covering the PG domain and test them in competition ELISA
- 10 with M75. According to the results, this was followed by a series of 6-12 aa oligopeptides. A major part of the PG domain consists of a 6-fold tandem repeat of 6 aa (aa 61 - 96) [SEQ ID NO: 97]; 4 repeats are identical (GEEDLP) [SEQ ID NO: 98] and 2 contain 2 aa exchanged (SEEDSP [SEQ ID NO: 141] and REEDPP [SEQ ID NO: 142]).

- Following are the results of competition ELISA with recombinant MN/CA
- 15 IX and oligopeptides synthesized according to partial sequences of the PG region. MN/CA IX+ and Δ produced in mammalian cells possessed a higher serological activity than any other protein or peptide included in this experiment; fusion protein GST-MN synthesized in bacteria was less active. The following peptides span the PG region: GGSSGEDDPLGEEDLPSEEDSPC (aa 51-72) [SEQ ID NO: 104];
- 20 GEEDLPSEEDSPREEDPPGEEDLPGEC (aa 61-85) [SEQ ID NO: 105]; EDPPGEEDLPGEEDLPGEEDLPVC (aa 75-98) [SEQ ID NO: 106]; and EVKPKSEEEGSLKLE (aa 97 - 111) [SEQ ID NO: 118]. SEQ ID NOS: 104 and 106 caused 50% inhibition at 1ng/ml. Those 2 oligopeptides are mutually non-overlapping, thus the epitope is repeated in both of them. SEQ ID NO: 105 was 1000x less active,
- 25 probably due to a different conformation. SEQ ID NO: 118 was inactive; thus it does not contain the M75 epitope.

- The next step for identifying the epitope was to synthesize oligopeptides containing all circular permutations of the motif GEEDLP [SEQ ID NO: 98] repeated twice. All 6 of the following dodecapeptides [SEQ ID NOS: 119-124] were
- 30 serologically active (2 more and 4 less so): GEEDLPGEEDLP [SEQ ID NO: 119]; EEDLPGEEDLP [SEQ ID NO: 120]; EDLPGEEDLP [SEQ ID NO: 121]; DLPGEEDLP [SEQ ID NO: 122]; LPGEEDLPGEED [SEQ ID NO: 123]; and PGEEDLPGEEDL [SEQ ID

NO: 124]. The following series of 7 aa sequences, flanked by alanine on both ends were tested: APGEEDLPA [SEQ ID NO: 125]; AGEEDLPGA [SEQ ID NO: 126]; AEEDLPGEA [SEQ ID NO: 127]; AEDLPGEEA [SEQ ID NO: 128]; ADLPGEEDA [SEQ ID NO: 129]; and ALPGEEDLA [SEQ ID NO: 130]. The results showed that the minimum serologically active sequence is the oligopeptide APGEEDLPA [SEQ ID NO: 125]. SEQ ID NOS: 127-130 proved negative in competition at 100 μ g/ml. Further, none of the following still shorter oligopeptides (6 + 2aa) competed in ELISA for M75: AGEEDLPA [SEQ ID NO: 131]; AEEDLPGA [SEQ ID NO: 132]; AEDLPGEA [SEQ ID NO: 133]; ADLPGEEA [SEQ ID NO: 134]; ALPGEEDA [SEQ ID NO: 135]; and APGEEDLA [SEQ ID NO: 136].

In the oligopeptides of SEQ ID NOS: 104, 105, 106 and 118, the C-terminal amino acid was present as an acid, whereas in all the other oligopeptides, the C-terminal amino acid was present as an amide. It is clear that the affinity between these oligopeptides and MAb M75 very strongly increases with the size of peptide molecule.

Attempts to demonstrate adhesion of cells to immobilized oligopeptides.

Our initial plan was to follow the pioneering work of Piersbacher and Ruoslahti, PNAS, 81: 5985 (1984). They linked tested oligopeptides to adsorbed bovine serum albumin by cross-linking agent SPDP (N-succinimidyl 3[pyridylhydro] propionate). This is why we added onto the C-end of oligopeptides SEQ ID NOS: 104-106 cysteine, which would enable oriented linking to adsorbed albumin. We demonstrated linking of the peptides directly in Petri dishes by immunoperoxidase staining with M75. Unfortunately, CGL1 or CGL3 cells adhered to control albumin treated with SPDP and blocked with ethanolamine (in place of oligopeptides) as strongly as to BSA dots with linked oligopeptides. We were unable to abrogate this non-specific adhesion. Oligopeptides SEQ ID NOS: 104-106 adsorb only very poorly to bacteriological Petri dishes, thereby not allowing the performance of the cell adhesion assay.

Alternatively, we tested inhibition of cell adhesion to MN/CA IX dots by oligopeptides added to the media together with the cell suspension, as described by Piersbacher and Ruoslahti, supra.. All peptides SEQ ID NOS: 104-106 and 118-136, were tested at concentrations of 100 and 10 μ g/ml. None of them inhibited reproducibly the adhesion of CGL1 cells.

Oligopeptides with affinity to M75 epitope which inhibit cell adhesion to MN/CA IX. As an alternative to monoclonal antibodies, we set out to select oligopeptides exerting affinity to M75 epitope as well as to MN/CA IX receptor binding site from a phage display library of random heptapeptides - Ph.D.-7. Our aim was to select phages containing the desired heptapeptides by panning on immobilized peptide SEQ ID NO: 106 and subsequent elution with M75. Eluted phage was multiplied in appropriate bacteria and subjected to 4 more cycles of panning and elution. From the selected phage population, 10 plaques were picked, amplified and the heptapeptide-coding region was sequenced. Only 3 heptapeptides were represented. Those three heptapeptides, after adding alanine on both sides, are the following nonapeptides: AKKMKRRKA [SEQ ID NO: 137]; AITFNAQYA [SEQ ID NO: 138]; and ASASAPVSA [SEQ ID NO: 139]. The last heptapeptide, synthesized again with added terminal alanines as nonapeptide AGQTRSPLA [SEQ ID NO: 140], was identified by panning on GST-MN and eluted with acetazolamide. This last peptide has affinity to the active site of MN/CA IX carbonic anhydrase. We synthesized these peptides of 7 + 2 aa and tested them in competition ELISA and in cell adhesion inhibition. Both tests yielded essentially consistent results: peptide SEQ ID NO: 138 showed the highest activity, peptide SEQ ID NO: 137 was less active, peptide SEQ ID NO: 139 was marginally positive only in ELISA, and peptide SEQ ID NO: 140 was inactive. In all of those 4 nonapeptides, the C-terminal amide was present as amide.

Discussion

Purification of transmembrane proteins like MN/CA IX often poses technical problems because they tend to form aggregates with other membrane proteins due to their hydrophobic TM segments. To avoid this, we engineered truncated MN/CA IX Δ IC Δ TM, which is secreted into the medium. Indeed, truncated MN/CA IX was obtained in higher purity than MN/CA IX+. Unfortunately, this protein was of little use for our purposes, since it was inactive in the cell adhesion assay. Such a situation has also been described for other cell adhesion molecules: their shed, shortened form either assumes an inactive conformation, or it adsorbs to hydrophobic plastic "upside down," while complete proteins adsorb by hydrophobic TM segments in the "correct" position.

MN/CA IX protein forms oligomers of 150 kDa, linked by disulfidic bonds. It was not known whether these are homo- or hetero-oligomers, but PAGE and Western blot analysis suggest that these are probably homo-oligomers, most likely trimers, since on the gel stained with Coomassie Blue no additional bands of intensity comparable to 2 bands specific for MN/CA IX appeared. It is also unlikely that there could exist an additional protein co-migrating with one of the 2 major MN/CA IX bands, since the intensity of their staining on the gel and on Western blots is well comparable.

There can be no doubt on the specificity of cell attachment to purified MN/CA IX+. It is abrogated by specific MAb M75, at a dilution 1:1000 of ascites fluid. This is a correction to our previous report in Zavada et al., Int. J. Oncol., 10: 857 (1997) in which we observed that MN/CA IX produced by vaccinia virus vector and fusion protein GST-MN support cell adhesion, but we did not realize that GST anchor itself contains another binding site, which is not blocked by M75.

MAb M75 reacts excellently with MN/CA IX under any circumstances - with native antigen on the surface of living cells, with denatured protein on Western blots and with antigen in paraffin sections of biopsies fixed with formaldehyde, suggesting that the epitope is small and contiguous. In competition ELISA the smallest sequence reactive with M75 was 7 + 2 aa, but the affinity between M75 and tested peptides strongly depended on their molecular weight. Complete MN/CA IX was 100,000x more active than the smallest serologically active peptide in terms of weight/volume concentration. In terms of molar concentration this difference would be 150,000,000x. Oligopeptides of intermediate size also showed intermediate activities. It remains to be elucidated whether such differences in activity are due to the conformation depending on the size of the molecule, or to the fact that complete MN/CA IX contains several copies of the epitope, but the smallest molecule only one.

Considering the possibility that the epitope is identical with the cell adhesion structure in MN/CA IX, we can understand why we failed to detect inhibition of cell adhesion by the oligopeptides. The binding site is just not as simple as the prototype peptide, RGD [Winter, J., IN Cleland and Craik (eds.), Protein Engineering. Principles and Practice, pp. 349-369 (N.Y.; Wiley-Liss; 1996)].

Naturally, one can argue that the size of MN/CA IX is about the same as of immunoglobulin molecule, and that binding of M75 to its epitope may sterically hinder

a different sequence of cell attachment site. This objection has been made unlikely by blocking of both M75 epitope and of cell binding site by nonapeptides 7 + 2 aa. That result strongly suggests that the epitope and the binding site are indeed identical.

MN/CA IX and its PG region in particular appears to be a potential target molecule for therapy for the following reasons: (i) it is exposed on the cell surface; (ii) it is present in high percentage of certain human carcinomas; (iii) it is normally expressed MN/CA IX in the mucosa of alimentary tract which is not accessible to circulating antibodies, in contrast with the tumors; (iv) it is not shed (or only minimally) into the body fluids; (v) the motif GEEDLP [SEQ ID NO: 98] is repeated 18 x on the surface of every MN/CA IX molecule. Oligopeptide display libraries are being employed in the first steps to develop new drugs [Winter, J., supra]. Selected oligopeptides can serve as lead compounds for the computerized design of new molecules, with additional properties required from a drug [DeCamp et al., IN Cleland and Craik (eds.), supra at pp. 467-505].

15

Example 3

Identification of Peptides Binding to MN Protein Using Phage Display

(a) To identify peptides that are recognized by MN protein, a heptapeptide phage display library [Ph.D.[®]-7 Peptide 7-mer Library Kit (phage display peptide library kit); New England Biolabs; Beverly, MA (USA)] was screened. In screening the library, a selection process, i.e., biopanning [Parmley and Smith, Gene, 73: 308 (1988); Noren, C.J., NEB Transcript, 8(1): 1 (1996)] was carried out by incubating the phages encoding the peptides with a plate coated with MN protein, washing away the unbound phage, eluting and amplifying the specifically bound phage.

The target MN protein in this process was a glutathione-S-transferase (GST) MN fusion protein (GST-MN). GST-MN is a recombinantly produced fusion protein expressed from pGEX-3X-MN containing the cDNA for the MN protein without the signal peptide. GST-MN was produced in bacteria under modified cultivation conditions (decreased optical density, decreased temperature). Such cultivation prevented premature termination of translation and resulted in synthesis of the protein molecules which were in vast majority of the full length. The GST-MN protein was

30

used for coating of the wells and binding the relevant phages. The bound phages were then eluted by acetazolamide, amplified and used for two additional rounds of screening.

After sequencing of several independent phage clones obtained after the third round of screening, the following heptapeptides were obtained:

(1) GETRAPL (SEQ ID NO: 107)

(2) GETREPL (SEQ ID NO: 108)

(3) GQTRSPL (SEQ ID NO: 109)

(4) GQTRSPL (")

(5) GQTRSPL (")

(6) GQTRSPL (")

(7) GQTRSPL (")

The heptapeptides show very similar or identical sequences indicating that the binding is specific. The fact that phages bearing these heptapeptides were eluted by acetazolamide, an inhibitor of carbonic anhydrase activity, indicates that the peptides bind to the CA domain of MN protein.

(b) Analogous screening of the heptapeptide phage display library is done using collagen I, shown to bind MN protein, for elution of phages. Different peptide(s) binding to different part(s) of the MN protein molecule are expected to be identified. After identifying such MN-binding peptides, the corresponding synthetic peptides shall then be analysed for their biological effects.

Example 4

Accessibility In Vivo of MN Protein Expressed in Tumor Cells and in Stomach

Lewis rats (384g) carrying a BP6 subcutaneous tumor (about 1 cm in diameter) expressing rat MN protein were injected intraperitoneally (i.p.) with ¹²⁵I-M75 Mab (2.5 x 10⁶ cpm). Five days later, 0.5-1g pieces of the tumor and organs were weighed and their radioactivity was measured by a gamma counter.

Table 2? summarizes the results. The highest radioactivity was present in the tumor. Relatively high radioactivity was found in the liver and kidney, apparently reflecting the clearance of mouse IgG from the blood. The stomach continued a

relatively low level of radioactivity, indicating that the M75 Mab had only limited access to MN protein exposed in the gastric mucosa.

TABLE 2

Distribution of radioactivity of ^{125}I -M75 in rat organs and in the tumor

	<u>Organ</u>	<u>cpm/g</u>				
10	Kidney	2153	2184			
	Spleen	653	555			
	Liver	1993	1880			
	Lung	1183	1025			
	Blood	1449				
15	Heart	568	477			
	Stomach	1184	1170			
	Testis	812	779			
	Tail	647				
	Tumor	3646	4058	3333	8653	3839

Example 5

FACS Analysis of MN Protein Expression in CGL3 Cells – Apoptosis

25 A FACS investigation was designed to determine the conditions that influence the synthesis of MN protein and to analyse the cell cycle distribution of MN-positive versus MN-negative cells in a CGL3 population stimulated to apoptosis. Previous Western blotting analyses have shown CGL3 cells to express a relatively high
30 amount of MN protein under different cultivation conditions. CGL3 cells are considered a constitutive producer of MN proteins. However, Western blotting does not recognize small differences in the level of protein. In contrast FACS allows the detection of individual MN-positive cells, a calculation of their percentage in the analysed population, an estimation of the level of MN protein in the cells, and a
35 determination of the cell cycle distribution.

To study the effect of cultivation conditions on MN expression in CGL3 cells, the CGL3 cells were plated in different relative densities and serum concentrations. Three days after plating, the cells were collected, surface labeled by M75 Mab followed by FITC-conjugated anti-mouse IgG and immediately analysed by FACS.

The analysis showed that in adherent cells, MN expression is dependent on cell density as is HeLa cells. However, low density cultures still produced detectable amounts of MN protein. In low density cultures, serum concentration does not seem to play a role. In relatively high density cultures, a decreasing serum concentration resulted in slightly diminished MN expression, probably due to a lower density that the cells were able to reach during the three days of cultivation.

The effect of the actual cell density is remarkable, and MN expression (detectable in 15-90% of the cells) represents a very sensitive monitoring factor. In all experiments, there was about a 5% higher percentage of cycling cells in the MN-positive part of the population, compared to the MN-negative part. That fact prompted the analysis of the cell cycle distribution of MN-positive CGL3 cells under unfavorable growth conditions, that is, after induction of apoptosis.

Apoptosis

CGL3 cells were stimulated to apoptotic death by several drugs, including cycloheximide, actinomycin D and dexamethasone. The FACS study showed that the onset of apoptosis is delayed in MN-positive cells suggesting a protective role of MN in this process. It was also observed that the induction of apoptosis resulted in the down-regulation of MN expression in a time-dependent manner. That same phenomenon was described for Bcl-2 anti-apoptotic protein, and there is existing opinion that the down-regulation of certain regulatory genes during apoptosis sensitizes the cells to undergo apoptotic death. To prove the role of MN in apoptosis, a similar study with cells transfected by MN cDNA is to be performed.

The preliminary results indicate the possible involvement of MN in the suppression of apoptosis. The recent view that tumors arise both as a consequence of increased proliferation and decreased cell death appears to be consistent with the association of the MN protein with tumors in vivo.

ATCC Deposits

The materials listed below were deposited with the American Type Culture Collection (ATCC) now at 10810 University Blvd., Manassus, Virginia 20110-2209 (USA). The deposits were made under the provisions of the Budapest Treaty on the International Recognition of Deposited Microorganisms for the Purposes of Patent Procedure and Regulations thereunder (Budapest Treaty). Maintenance of a viable culture is assured for thirty years from the date of deposit. The hybridomas and plasmids will be made available by the ATCC under the terms of the Budapest Treaty, and subject to an agreement between the Applicants and the ATCC which assures unrestricted availability of the deposited hybridomas and plasmids to the public upon the granting of patent from the instant application. Availability of the deposited strain is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any Government in accordance with its patent laws.

15	<u>Hybridoma</u>	<u>Deposit Date</u>	<u>ATCC #</u>
	VU-M75	September 17, 1992	HB 11128
	MN 12.2.2	June 9, 1994	HB 11647
	<u>Plasmid</u>	<u>Deposit Date</u>	<u>ATCC #</u>
20	A4a	June 6, 1995	97199
	XE1	June 6, 1995	97200
	XE3	June 6, 1995	97198

The description of the foregoing embodiments of the invention have been presented for purposes of illustration and description. They are not intended to be exhaustive or to limit the invention to the precise form disclosed, and obviously many modifications and variations are possible in light of the above teachings. The embodiments were chosen and described in order to explain the principles of the invention and its practical application to enable thereby others skilled in the art to utilize the invention in various embodiments and with various modifications as are suited to the particular use contemplated.

All references cited herein are hereby incorporated by reference.

CLAIMS

1. An organic or inorganic molecule that binds specifically to a site on MN protein to which vertebrate cells adhere in a cell adhesion assay, wherein said molecule when tested in vitro inhibits the adhesion of cells to MN protein.
2. The organic or inorganic molecule of Claim 1 wherein said molecule, when in contact with a vertebrate preneoplastic or neoplastic cell that abnormally expresses MN protein, inhibits the growth of said cell.
3. The molecule of Claims 1 and 2 which is organic.
4. The molecule of Claim 3 which is a protein or a polypeptide.
5. The molecule of Claim 4 wherein said protein or polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 107, 108, 109, 137 and 138.
6. The molecule of Claim 4 wherein said polypeptide is selected from the group consisting of SEQ ID NOS: 107, 108, 109, 137 and 138.
7. The molecule of Claim 1 wherein the site on the MN protein to which said vertebrate cells adhere in said cell adhesion assay is within the proteoglycan-like domain or within the carbonic anhydrase domain of the MN protein.
8. The molecule of Claim 1 wherein the site on the MN protein comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 10 and 97-106.
9. The molecule of Claim 1 wherein the site on the MN protein has an amino acid sequence selected from the group consisting of SEQ ID NOS: 10 and 97-106.

10. The molecule of Claim 1 wherein said vertebrate cells are mammalian.

11. The molecule of Claim 1 wherein said vertebrate cells are human.

12. A MN protein or MN polypeptide which mediates attachment of vertebrate cells in a cell adhesion assay, wherein said MN protein or MN polypeptide when introduced into the extracellular fluid environment of vertebrate cells prevents the formation of intercellular contacts and the adhesion of said vertebrate cells to each other.

13. The MN protein or MN polypeptide of Claim 12 which when introduced into the extracellular fluid environment of a vertebrate preneoplastic or neoplastic cell that abnormally expresses MN protein, inhibits the growth of said cell.

14. The MN protein or MN polypeptide of Claim 12 whose amino acid sequence is from SEQ ID NO: 97, from SEQ ID NO: 50, or from SEQ ID NO: 51.

15. The MN protein or MN polypeptide of Claim 12 whose amino acid sequence is from SEQ ID NO: 50.

16. The MN protein or MN polypeptide of Claim 12 which comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 10 and 97-106.

17. The MN polypeptide of Claim 12 that is selected from the group consisting of SEQ ID NOS: 10 and 97-106.

18. The MN protein or MN polypeptide of Claim 12 that is specifically bound by either the M75 monoclonal antibody that is secreted from the hybridoma VU-M75, which was deposited at the American Type Culture Collection under ATCC No. HB 11128, or by the MN12 monoclonal antibody that is secreted from the hybridoma

MN 12.2.2, which was deposited at the American Type Culture Collection under ATCC No. HB 11647, or by both said monoclonal antibodies.

19. A method of identifying a site on an MN protein to which vertebrate
5 cells adhere by testing a series of overlapping polypeptides from said MN protein in a cell adhesion assay with vertebrate cells, and determining that if cells adhere to a polypeptide from said series, that said polypeptide comprises a site on said MN protein to which vertebrate cells adhere.

10 20. A vector comprising an expression control sequence operatively linked to a nucleic acid encoding the variable domains of a MN-specific antibody, wherein said domains are separated by a flexible linker polypeptide, and wherein said vector, when transfected into a vertebrate preneoplastic or neoplastic cell that abnormally expresses MN protein, inhibits the growth of said cell.

15 21. The vector of Claim 20 wherein said expression control sequence comprises the MN gene promoter operatively linked to said nucleic acid.

22. The vector of Claim 20 wherein said flexible linker polypeptide has
20 the amino acid sequence of SEQ ID NO: 116.

23. The vector of Claim 20 wherein said expression control sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 27 and SEQ ID NO: 91.

25 24. A vector comprising a nucleic acid that encodes a cytotoxic protein or cytotoxic polypeptide operatively linked to the MN gene promoter, wherein said vector, when transfected into a vertebrate preneoplastic or neoplastic cell that abnormally expresses MN protein, inhibits the growth of said cell.

30 25. The vector of Claim 24 wherein said cytotoxic protein is HSV thymidine kinase.

26. The vector according to Claim 24 wherein said vector further comprises a nucleic acid encoding a cytokine operatively linked to said MN gene promoter.

5 27. The vector of Claim 26 wherein said cytokine is interferon or interleukin-2.

28. A repressor complex that binds to the MN gene promoter.

10 29. The repressor complex of Claim 28 that binds to SEQ ID NO: 115 of the MN gene promoter.

30. The repressor complex of Claim 29 comprising two proteins having molecular weights of 35 and 42 kilodaltons, respectively.

15

1	1	ACA	GTC	AGC	CGC	ATG	GCT	CCC	CTG	TGC	CCC	AGC	CCC	TGG	CTC	CCT	CTG	12	48
13	49	L	I	P	A	P	A	P	G	L	T	V	Q	L	L	L	S	28	96
29	97	L	L	L	L	M	P	P	V	H	P	Q	R	L	P	R	M	44	144
45	145	E	D	S	P	L	G	G	G	S	S	G	E	D	D	P	L	60	192
61	193	G	E	E	D	L	P	S	E	E	D	S	P	R	E	E	D	76	240
77	241	P	P	G	E	E	D	L	P	G	E	E	D	L	P	G	E	92	288
93	289	E	D	L	P	E	V	K	P	K	S	E	E	E	G	S	L	108	336
109	337	K	L	E	D	L	P	T	V	E	A	P	G	D	P	Q	E	124	384
125	385	P	Q	N	N	A	H	R	D	K	E	G	D	D	Q	S	H	140	432
141	433	W	R	Y	G	G	D	P	P	W	P	R	V	S	P	A	C	156	480
157	481	A	G	R	F	Q	S	P	V	D	I	R	P	Q	L	A	A	172	528

FIG.-1A

2 / 13

```

173 F C P A L R P L E L L G L Q L P 188
529 TTC TGC CCG GCC CTG CTG CGC CCC CTG GAA CTC CTG GGC TTC CAG CTC CCG 576

189 P L P E L R L R N N G H S V Q L 204
577 CCG CTC CCA GAA CTG CTG CGC CTG CGC AAC AAT GGC CAC AGT GTG CAA CTG 624

205 T L P P G L E M A L G P G R E Y 220
625 ACC CTG CCT CCT GGG CTA GAG ATG GCT CTG GGT CCC GGG CGG GAG TAC 672

221 R A L Q L L H L H W G A A G R P G 236
673 CGG GCT CTG CAG CTG CAT CTG CAC TGG GGG GCT GCA GGT CGT CCG GGC 720

237 S E H T V E G H R F P A E I H V 252
721 TCG GAG CAC ACT GTG GAA GGC CAC CGT TTC CCT GCC GAG ATC CAC GTG 768

253 V H L S T A F A R V D E A L G R 268
769 GTT CAC CTC AGC ACC GCC TTT GCC AGA GTT GAC GAG GCC TTG GGG CGC 816

269 P G G L A V L A A F L E E G P E 284
817 CCG GGA GGC CTG GCC GTG TTG GCC TTT CTG GAG GAG GGC CCG GAA 864

285 E N S A Y E Q L L S R L E E I A 300
865 GAA AAC AGT GCC TAT GAG CAG TTG CTG TCT TCT CGC TTG GAA GAA ATC GCT 912

301 E E G S E T Q V P G L D I S A L 316
913 GAG GAA GGC TCA GAG ACT CAG GTG GTC CCA GGA CTG GAC ATA TCT GCA CTC 960

317 L P S D F S R Y Y F Q Y E G S L T 332
961 CTG CCC TCT GAC TTC AGC CGC TAC TTC CAA TAT GAG GGG TCT CTG ACT 1008

333 T P P C A Q G V I W T V F N Q T 348
1009 ACA CCG CCC TGT GCC CAG GGT GTC ATC TGG ACT GTG TTT AAC CAG ACA 1056

```

FIG.-1B

349 V M L S A K Q L H T L S D T L W 364
1057 GTG ATG CTG AGT GCT AAG CAG CTC CAC ACC CTC TCT GAC ACC CTG TGG 1104

365 G P G D S R L Q L N F R A T Q P 380
1105 GGA CCT GGT GAC TCT CGG CTA CAG CTG AAC TTC CGA GCG ACG CAG CCT 1152

381 L N G R V I E A S F P A G V D S 396
1153 TTG AAT GGG CGA GTG ATT GAG GCC TCC TTC CCT GCT GGA GTG GAC AGC 1200

397 S P R A A E P V Q L N S C L A A 412
1201 AGT CCT CGG GCT GCT GAG CCA GTC CAG CTG AAT TCC TGC CTG GCT GCT 1248

413 G D I L A L V F G L L F A V T S 428
1249 GGT GAC ATC CTA GCC CTG GTT TTT GGC CTC CTT TTT GCT GTC ACC AGC 1296

429 V A F L V Q M R R Q H R R G T K 444
1297 GTC GCG TTC CTT GTG CAG ATG AGA AGG CAG CAC AGA AGG GGA ACC AAA 1344

445 G G V S Y R P A E V A E T G A * 460
1345 GGG GGT GTG AGC TAC CGC CCA GCA GAG GTA GCC GAG ACT GGA GCC TAG 1392

1393 AGG CTG GAT CTT GGA GAA TGT GAG AAG CCA GCC AGA GGC ATC TGA GGG 1440

1441 GGA GCC GGT AAC TGT CCT GTC CTG CTC ATT ATG CCA CTT CCT TTT AAC 1488

1489 TGC CAA GAA ATT TTT TAA AAT AAA TAT TTA TAA T 1522

FIG.-1C**FIG.-1**

FIG.-1A

FIG.-1B

FIG.-1C

```

1  ggatcctgtt gactcgtgac ctaccccca accctgtgct ctctgaaca tgagctgtgt
61  ccactcaggg ttaaatggat taaggcggt gcaagatgtg ctttgttaaa cagatgcttg
121 aaggcagcat gctcgttaag agtcatcacc aatccctaata ctaagtaat cagggacaca
181 aacactgcgg aaggccgcag ggtcctctgc ctattgtcca tgaccctgcc aaatccccct ctgtgagaaa
241 tttatctgac cttccctcca aaaaataaat taaaaaaa atacaaaaa aaaaaaaaa
301 cacccaagaa ttatcaataa gacttacgaa tagttattga taaatgaata gctattggta aagccaagta
361 aaaaaaaaaa gacttacgaa gacggccatc gacggccatc atcacagctc aagtctacct gatttgatct
421 aatgatcata ttcaaaacca gacggccatc gatttcacta gattagtcac catcctcaaa attctcccc
481 ctttatcatt gtcattcttt gatttcacta gattagtcac gattagtcac catcctcaaa attctcccc
541 aagtttctaat tacgtttccaa acatttaggg gttacatgaa gttgaacct actaccttct
601 ttgcttttga gccatgagtt gtaggaatga gcttttggt tatttttgta gctaattttg
661 tttaaaacttt acctctaagt cagttgggta gcctttggct tatttttgta gctaattttg
721 tagttaatgg atgcactgtg aatcttgcta tgatagtttt cctccacct ttgccactag
781 gggtaggtag gtactcagtt ttccagtaatt gcttacctaa gaccctaagc cctattttctc
841 ttgtactggc ctttatctgt aatatgggca tatttaatac aatataattt ttggagtttt
901 ttgtttgtt tgttgtttg ttttttgag acggagtctt gcatctgtca tgcccaggct
961 ggagtagcag tggtgccatc tcggctcact gcaagctcca cctcccgagt tcacgccatt
1021 ttcctgcctc agcctccga gtagctggga ctacaggcg cggccacct gccgggctaa
1081 ttttttgtat ttttggtaga gacggggttt caccgtgtta gccagaatgg tctcgatctc
1141 ctgacttcgt gatccacccg cctcggcctc ccaaagtctt gggattacag gtgtgagcca
1201 ccgcacctgg ccaattttt gagtctttta aagtaaaaat atgtcttcta agctggtaac
1261 tatggtagat ttctttttat taatgtgggtg ctgacgggtca tataggttctt tttgagtttg
1321 gcatgcatat gctacttttt gcagtccttt cattacattt ttctctcttc atttgaagag
1381 catgtttatat ctttttagctt cacttggctt aaaagggttct ctattagcc taacacagtg
1441 tcattgttgg taccacttgg atcataagtg gaaaaacagt caagaaatg cacagtaata
1501 cttgttttga agaggatga ttcagggtgaa tctgacacta agaaactccc ctacctgagg
1561 tctgagattc ctctgacatt gctgtatata ggcttttctt ttgacagcct gtgactgcgg
1621 actatttttc ttaagcaaga tatgctaag tatgttgagc ctttttccag agagagggtct
1681 catatctgca tcaagtgaga acatataatg tctgcatgtt tccatatttc aggaatgttt
1741 gcttgtgttt tatgctttta tatagacagg gaaacttgtt cctcagtgac ccaaaagagg
1801 tgggaattgt tattggatat catcattggc ccacgctttc tgaccttgga aacaattaa
1861 ggttcataat ctcaattctg tcagaattgg tacaagaaat agctgctatg tttcttgaca
1921 tccacttgg taggaaataa gaatgtgaaa ctcttcagtt ggtgtgtgtc cct?gtttt

```

FIG._2A

5 / 13

1981 ttgcaatttc ctcttactg tgttaaaaa aagtatgatc ttgctctgag aggtgaggca
2041 ttcttaatac tgatctttaa agatcaataa tataatcctt tcaaggatta tgtctttatt
2101 ataataaaga taatttgtct ttaacagaat caataatata atcccttaa ggattatatc
2161 ttgctgggc gcagtggctc acacctgtaa tcccagcact ttgggtggcc aaggtggaag
2221 gatcaaatct gcctacttct atattatctt ctaaagcaga attcatctct ctccctcaa
2281 tatgatgata ttgacagggt ttgacctcac tccactagatt gtgagctcct gctcagggca
2341 ggtagcgttt ttgttttttg tttttgtttt tcttttttga gacagggtct tgctctgtca
2401 ccaggccag agtgcaatgg tacagtctca gctcactgca gcctcaacc cctcggctca
2461 aaccatcatc ccatctcagc ctccctgagta gctgggacta caggcacatg ccattacacc
2521 tggctaattt ttttgtattt ctagttaga cagggtttgg ccatgttgc cgggctggtc
2581 tcgaactcct ggactcaagc aatccaccca cctcagcctc ccaaatgag ggaccgtgtc
2641 ttattcatct ccatgtccct agtccatagc ccagtgtgg acctatgga gtactaaata
2701 aatatctgtt gaatgcaata gtaaatagca ttccaggag caagaactag ataacaaaag
2761 gtggtaaaag gtttgagaa aaaaataata gtttaatttg gctagagtat gaggagagt
2821 agtaggagac aagatggaaa ggtctcttgg gcaaggtttt gaaggaagtt ggaagtcaga
2881 agtacacaat gtgcatactg tggcaggcag tggggagcca atgaaggctt ttgagcagga
2941 gagtaatgtg ttgaaaaata aatataggtt aaacctata gagccctct gacacataca
3001 ctgtctttc attcaagctc aagtttgtct ccacatacc cattactaa ctacccctg
3061 ggctccctta gcagcctgcc ctacctctt ctacctctt accctgttc tggaggatc agggatgtat
3121 acatgagctg cttccctct cagccagagg acatgggggg cccagctcc cctgccttcc
3181 cccttctgtg cctggagctg ggaagcaggc cagggttagc tgaggctggc tggcaagcag
3241 ctgggtgggt ccaggagag cctgcatagt gccagggtgt gccttgggtt ccaagctagt
3301 ccatggcccc gataacctc tgcctgtgca cacacctgcc cctcactca ccccatcct
3361 agctttggta tgggggagag ggcacagggc cagacaaacc tgtgagactt tggctccatc
3421 tctgcaaaag ggcgtctgt gagtacgctt gctccctcc aggttgtct cccccacc
3481 cagctctcgt ttccaatgca cgtacagccc gtacacaccg tgtgctggga caccacacag
3541 TCAGCCGCAT GGCTCCCTG TGCCCCAGCC CCTGGCTCCC TCTGTTGATC CCGGCCCTG
3601 CTCAGGCCT CACTGTGCAA CTGCTGCTGT CACTGCTGCT TCTGGTGCCT GTCCATCCCC
3661 AGAGGTTGCC CCGGATGCAG GAGGATTCCC CTTGGGAGG AGGCTCTTCT GGGGAAGATG
3721 ACCCACTGG CGAGGAGGAT CTGCCCAGTG AAGAGGATTC ACCAGAGAG GAGGATCCAC
3781 CCGGAGAGGA GGATCTACCT GGAGAGGAGG ATCTACCTGG AGAGGAGGAT CTACCTGAAG
3841 TTAAGCCTAA ATCAGAAGAA GAGGGCTCCC TGAAGTTAGA GGATCTACCT ACTGTTGAGG
3901 CTCTGGAGA TCCTCAAGAA CCCAGAATA ATGCCACAG GGACAAAGAA Gtaagtgg

FIG.-2B

3961 catcaatctc caaatccagg ttccaggagg ttcatgactc cctcccata cccagccta
4021 ggctctgttc actcagggaa ggaggggaga ctgtactccc cacagaagcc cttccagagg
4081 tccataacca atatcccat cccactctc ggaggtagaa agggacagat gtggagagaa
4141 aataaaaaagg gtgcaaaaagg agagaggtga gctggatgag atgggagaga agggggaggc
4201 tggagaagag aaaggatga gaactgcaga tgagagaaaa aatgtgcaga cagaggaaaa
4261 aaataggtgg agaaggagag tcagagagtt tgaggggaag agaaaaaggaa agcttgggag
4321 gtgaagtggg taccagagac aagcaagaag agctggtaga agtcatctca tcttaggcta
4381 caatgaggaa ttgagaccta ggaagaaggg acacagcagg tagagaaaacg tggcttcttg
4441 actcccaagc caggaatttg gggaaaagggg ttggagacca tacaaggcag agggatgagt
4501 ggggagaaga aagaaggag aaaggaaga tggtgtactc actcatttgg gactcaggac
4561 tgaagtggc actcactttt tttttttttt tttttgagac aaactttcac ttttgttgcc
4621 caggctggag tgcaatggcg cgatctcggc tactgcaac ctccacctcc cgggttcaag
4681 tgattctcct gcctcagcct ctagccaagt agctgcgatt acaggcatgc gccaccacgc
4741 cgggctaatt tttgtatttt tagtagagac ggggtttcgc catgttggtc aggtgggtct
4801 cgaactcctg atctcaggtg atccaaccac cctggcctcc caaagtgtg ggaattatagg
4861 cgtgagccac aggcctggc ctgaagcagc cactcacttt tacagaccct aagacaatga
4921 ttgcaagctg gtaggattgc tgtttggccc accagctgc ggtgttagt ttgggtgcgg
4981 tctcctgtgc ttgacactg gccgcctaa ggcatttgtt accgtaatg ctctgtgaag
5041 gcatctgcgt ttgtgacatc gttttggtcg ccaggaaggg attggggctc taagcttgag
5101 cggttcatcc ttttcattta tacagGGGAT GACCAGAGTC ATTGGCGCTA TGGAGgtgag
5161 acaccacccc gctgcacaga cccaatctgg gaaccagct ctgtggatct cccctacagc
5221 cgtccctgaa cactggtccc gggcgtccca ccgcccgcct accgtcccac cccctcacct
5281 ttcttaccg ggttccctaa gttcctgacc taggcgtcag acttcctcac tatactctcc
5341 caccacagc GACCCGCCCT GGCCTCCGGT GTCCCCAGCC TGCCTGGCC GCTTCCAGTC
5401 CCCGGTGGAT ATCCGCCCCC AGCTCGCCCG CTTCCTGCCG GCCCTGCCG CCTGGAACCT
5461 CCTGGGCTTC CAGCTCCCCG CGTCCCAGA ACTGCGCCTG GCACAATG GCCACAGTGg
5521 tgagggggtc tccccgcga gacttgggga tggggcgggg cgcagggaag ggaaccgtcg
5581 cgcagtgcct gcccggggt tgggctggcc ctaccgggcg gggccggctc acttgctct
5641 ccctacgcag TGCAACTGAC CCTGCCCTCCT GGGCTAGAGA TGGCTCTGGG TCCCGGCGCG
5701 GAGTACCGGG CTCTGCAGCT GCATCTGCAC TGGGGGGCTG CAGGTCGTCC GGGCTCGGAG
5761 CACACTGTGG AAGGCCACCG TTTCCCTGCC GAGgtgagcg cggactggcc gagaaggggc
5821 aaaggagcgg ggcggacggg ggccagagac gtggccctct cctaccctcg tgtcctttc
5881 agATCCACGT GGTTCACCTC AGCACCGCCT TTGCCAGAGT TGACGAGGCC TTGGGGCGCC

FIG.-2C

```

5941 CGGGAGGCCCT GCGCGTGTTG GCGGCCTTTC TGGAGgtacc agatcctgga cacccttac
6001 tccccgcttt cccatcccat gctcctcccg gactctatcg tggagccaga gaccctatcc
6061 cagcaagctc actcaggccc ctggctgaca aactcattca cgcactgttt gtccatttaa
6121 caccactgt gaaccaggca ccagccccc acaaggattc tgaagctgta ggtccttgcc
6181 tctaaggagc ccacagccag tgggggaggg tgacatgaca gacacatagg aaggacatag
6241 taaagatggt ggtcacagag gaggtgacac ttaaagcctt cactggtaga aaagaaaagg
6301 aggtgttcat tgcagaggaa acagaaatgt caaagactca gaatatggcc tatttaggga
6361 atggctacat acaccatgat tagaggaggc ccagtaaaagg gaagggatgg tgagatgcct
6421 gctaggttca ctcactcact ttattattt tattatttt ttgacagtc tctctgtgc
6481 ccaggctgga gtgcagtgggt gtgatcttgg gtcactgcaa cttccgcctc ccgggttcaa
6541 gggattctcc tgcctcagct tcctgagtag ctggggttac aggtgtgtgc caccatgccc
6601 agctaatttt tttttgtatt tttagtagac aggtttcac catgttggtc aggtggtct
6661 caaactcctg gcctcaagtg atccgcctga ctcagcctac caaagtgtc attacaaagt
6721 tgagccaccg tgcccagcca cactcactga ttctttaatg ccagccacac agcacaaagt
6781 tcagagaaat gcctccatca tagcatgtca atatgttcat actcttaggt tcatgatgtt
6841 cttaacatta ggttcataag caaaataaga aaaaagaata ataaataaaa gaagtggcat
6901 gtcaggacct cacctgaaa gctgataat gcttccctgt ggtgaatgca gaggtgacac
6961 caacacaaag gtgtatata cctcagaagg gcttccctgt ggggagtatg tacggaggca gcagtgaagt
7021 agactgcaaa cgtcagaagg gctcaggtca ctgagagcct agtatcctag taaagtggc
7081 tcttccctc tcttccagc ttgtcattga aaaccagtcc accaagcttg ttggttcgca
7141 cagcaagagt acatagagtt tgaataata taaggattt taagaggag acactgtctc
7201 taaaaaaaaa aacaacagca acaacaaaa ccactaatct gttcataaa aatcacctt
7261 agcatctca gagctgagga atgggagagg actatgggaa ccccttcat gtccggcct
7321 tcagccatgg ccctggatac atgcactcat ctgtcttaca atgtcattcc ccagGAGGG
7381 CCCGGAAGAA AACAGTGCCCT ATGAGCAGTT GCTGTCTCGC TTGGAAGAAA TCGCTGAGGA
7441 AGgtcagttt gttggtctgg ccactaatct ctgtggccta gttcataaa aatcacctt
7501 tggagcttca ggtctgaggg tggagatggg ctccctccag tgcaggaggg attgaagcat
7561 gagccagcgc tcatcttgat aataaccatg aagctgacag acacagttac ccgcaaacgg
7621 ctgcctacag attgaaaacc aagcaaaaac cgcggggcac ggtggctcac gcctgtaac
7681 ccagcacttt gggaggccaa ggcatgtgga tcacgaggtc aagagatcaa gaccatcctg
7741 gccaacatgg tgaaccccca tctctactaa aaatacgaaa aaatagccag gcgtggtggc
7801 ggggtgcctgt aatcccagct actcgggagg ctgaggcagg agaattggcat gaacccggga
7861 ggcagaagtt gcagtgaacc gagatcgtgc cactgcactc cagcctgggc aacagagcga

```

FIG._2D

7921 gactcttgct tcaaaaaaa aaaaaaaa gaaaccaag caaaaccaa aatgagacaa
7981 aaaaaacaag accaaaaaat ggtgtttgga aattgtcaag gtcaagtctg gagagctaaa
8041 ctttttctga gaactgttta tcgttctctt cttagtcact cttggtcat tttaaatctc acttactcta
8101 ttgttggaat taggtttctg cttagtcact cttagtcact ctttgcat tcttggtgtct
8161 cttagacctt taggtttctg cttagtcact cttagtcact ctttgcat tcttggtgtct
8221 gtttttgtata gttatcaata tttttttttt ttttttttat ttacaaagt tttaaatctc
8281 tctttttttt tctcaaaact tctcaaaact ctgacctgt gatccaccag cctcgccctc
8341 gccaggctgc tttttttttt tttttttttt aatttgctct gggcttaaac ttgtggccca
8401 gggattcatt tttttttttt tttttttttt tgtagactca gacgggtctt tcttctcct
8461 atggtacaca gagttaagag tttttttttt tttttttttt tttttttttt tcttctcct
8521 cctcccttcc cttccacctt caagcctct aagtggtct agagttgagt tcttggtgga
8581 caggcctctt ctagtgctc ctagtgctc aagtggtct agagttgagt tcttggtgga
8641 agggcctgca ctagtgctc ctagtgctc aagtggtct agagttgagt tcttggtgga
8701 gaaactgtat cctataccc tgaagcttta aggggtgca atgtagatga gacccaaca
8761 tagatcctct tcacagGCTC AGAGACTCAG GTCCACGGAC TGGACATATC TGCACCTCTG
8821 CCTCTGACT TCAGCCGCTA CTTCCAATAT GAGGGTCTC TGACTACACC GCCCTGTGCC
8881 CAGGGTGTCA TCTGGACTGT GTTTAACAG ACAGTGATGC TGAGTGCTAA GCAGgtgggc
8941 ctggggtgtg tgtggacaca gtgggtgctg gggaaagagg atgtaagatg agatgagaaa
9001 caggagaaga aagaaatcaa ggctgggctc tgtggcttac gcctataatc ccaccacgtt
9061 gggaggctga ggtgggagaa tggtttgagc ccaggagttc aagacaaggc ggggcaacat
9121 agtgtgacct catctctacc tactcaagga ggctgaggtg ggaagatcgc ttgattccag
9181 gtatgcggcc tagtcccagc ctatgatccc accactgcct accatctta ggatacat
9241 gagtttgaga ctgcagtga ctgcagtga aagaggctgg atggggaata caggagctgg
9301 atttatttat aaaagaaatc aagaggctgg atggggaata caggagctgg aggggtggagc
9361 cctgaggtgc tggttgtgag ctggcctggg acccttggtt cctgtcatgc catgaacca
9421 ccacactgt cactgacct ccctagCTCC ACACCTCTC TGACACCTCTG TGGGACCTG
9481 GTGACTCTCG GCTACAGCTG AACTTCCGAG CGACGCAGCC TTTGAATGGG CGAGTGATTG
9541 AGGCTCCTT CCTGCTGGA GTGGACAGCA GTCCTCGGC TGCTGAGCCA Ggtacagctt
9601 tgtctggtt ccccccagcc agtagtccc tatcctccc tcttggtgct agtgtctgtc
9661 attggtggtc acagccgccc tctcacatct cctttttctc tccagTCCAG CTGAATTCTT
9721 GCCTGGCTGC TGgtgagtct gcccctctc ttggtcctga tggcaggaga ctcctcagca
9781 ccattcagcc ccagggtgc tcaggacctg ccttgctccc tctccttctc tgcagaacag
9841 accccaacc caatataga gaggcagatc atggtgggga tcccccat gtccccagag

FIG._2E

9901 gctaattgat tagaatgaag cttgagaaat ctccagcat ccctctcgca aaagaatccc
 9961 cccccctttt tttaaagata gggctctcact ctgtttgccc caggctgggg tggtgtggca
 10021 cgatcatagc tcatgcagc ctgaaactcc taggtcagg caatcctttc accttagctt
 10081 ctcaaagcac tgggactgta ggcactgagc actgtgcctg gccccaaacg gcccttttac
 10141 ttggcctttta ggaagcaaaa acggtgctta tcttaccctt tctcgtgtat ccaccctcat
 10201 cccttggtg cctcttctg gagactgagg cactatgggg ctgcctgaga actcggggca
 10261 ggggtggtgg agtgactga ggcagggtgtt gaggaactct gcagacccct cttccttccc
 10321 aaagcagccc tctctgctct ccatcgcagg TGACATCCTA GCCCTGGTTT TTGGCCCTCCT
 10381 TTTTGTGTC ACCAGCGTCG CGTTCCTTGT GCAGATGAGA AGCAGCACCA Ggtattacac
 10441 tgaccctttc ttcagggcaca agcttcccc acccttgttg agtcacttca tgcaaaagcg
 10501 atgcaaatga gctgctcctg ggccagtctt ctgattagcc ttccctgttg tgtacacaca
 10561 GAAGGGGAAC CAAAGGGGT GTGAGCTACC GCCCAGCAGA GGTAGCCGAG ACTGGAGCCT
 10621 AGAGGCTGGA TCTTGGAGAA TGTGAGAAGC CAGCCAGAGG CATCTGAGGG GGAGCCGGTA
 10681 ACTGTCCTGT CCTGCTCATT ATGCCACTTC CTTTAACTG CCAAGAAATT TTTTAAATA
 10741 AATATTATA ATaaatatg tgtagtcac ctttgttccc caaatcagaa ggaggtattt
 10801 gaatttccta ttactgttat tagcaccaat ttagtggtaa tgcatttatt ctattacagt
 10861 tcggcctcct tccacacatc actccaatgt gttgctcc

FIG._2F

FIG._2A

FIG._2B

FIG._2C




FIG._2D

FIG._2E

FIG._2F

FIG._2

10 / 13

**FIG._3****FIG._4****FIG._8**




FIG._5

12 / 13

```

-506 CTTGCTTTTC ATTCAAGCTC AGTTTGCTCT CCCACATACC CATTAUCTTAA CTCACCCCTCG
-446 GGCTCCCCTA GCAGCCTGCC CTACCTCTTT ACCTGCTTCC TGGTGGAGTC AGGGATGTAT
      AP2
-386 ACATGAGCTG CTTTCCCTCT CAGCCAGAGG ACATGGGGGG CCCCAGCTCC CCTGCCTTTC
-326 CCCTTCTGTG CCTGGAGCTG GGAAGCAGGC CAGGGTTAGC TGAGGCTGGC TGGCAAGCAG
-266 CTGGGTGGTG CCAGGGAGAG CCTGCATAGT GCCAGGTGGT GCCTTGGGTT CCAAGCTAGT
      VII p53
-206 CCATGGCCCC GATAACCTTC TGCCTGTGCA CACACCTGCC CCTCACTCCA CCCCATCCT
      VI Inr V
-146 AGCTTTGGTA TGGGGGAGAG GGCACAGGGC CAGACAAACC TGTGAGACTT TGGCTCCATC
      IV AP1 III Inr
-86 TCTGCAAAAG GCGCTCTGT GAGTCAGCCT GCTCCCCCTCC AGGCTTGCTC CTCCCCCACC
      II AP1 I AP2
      ***
-26 CAGCTCTCGT TTCCAATGCA CGTACAGCCC GTACACACCG TGTGCTGGGA CACCCACAG
      ...

```

FIG._6




FIG. 7

SEQUENCE LISTING

<110> Zavada, Jan
 Pastorekova, Silvia
 Pastorek, Jaromir

<120> MN Gene and Protein

<130> D-0021.5 PCT

<140>

<141>

<150> 09/177,776

<151> 1998-10-23

<150> 09/178,115

<151> 1998-10-23

<160> 143

<170> PatentIn Ver. 2.0

<210> 1

<211> 1522

<212> DNA

<213> HUMAN

<220>

<221> CDS

<222> (13)..(1389)

<220>

<221> mat_peptide

<222> (124)..(1389)

<400> 1

acagtcagcc gc atg gct ccc ctg tgc ccc agc ccc tgg ctc cct ctg ttg 51
 Met Ala Pro Leu Cys Pro Ser Pro Trp Leu Pro Leu Leu
 -35 -30 -25

atc ccg gcc cct gct cca ggc ctc act gtg caa ctg ctg ctg tca ctg 99
 Ile Pro Ala Pro Ala Pro Gly Leu Thr Val Gln Leu Leu Ser Leu
 -20 -15 -10

ctg ctt ctg atg cct gtc cat ccc cag agg ttg ccc cgg atg cag gag 147
 Leu Leu Leu Met Pro Val His Pro Gln Arg Leu Pro Arg Met Gln Glu
 -5 -1 1 5

gat tcc ccc ttg gga gga ggc tct tct ggg gaa gat gac cca ctg ggc	195
Asp Ser Pro Leu Gly Gly Gly Ser Ser Gly Glu Asp Asp Pro Leu Gly	
10 15 20	
 gag gag gat ctg ccc agt gaa gag gat tca ccc aga gag gag gat cca	243
Glu Glu Asp Leu Pro Ser Glu Glu Asp Ser Pro Arg Glu Glu Asp Pro	
25 30 35 40	
 ccc gga gag gag gat cta cct gga gag gag gat cta cct gga gag gag	291
Pro Gly Glu Glu Asp Leu Pro Gly Glu Glu Asp Leu Pro Gly Glu Glu	
45 50 55	
 gat cta cct gaa gtt aag cct aaa tca gaa gaa gag ggc tcc ctg aag	339
Asp Leu Pro Glu Val Lys Pro Lys Ser Glu Glu Glu Gly Ser Leu Lys	
60 65 70	
 tta gag gat cta cct act gtt gag gct cct gga gat cct caa gaa ccc	387
Leu Glu Asp Leu Pro Thr Val Glu Ala Pro Gly Asp Pro Gln Glu Pro	
75 80 85	
 cag aat aat gcc cac agg gac aaa gaa ggg gat gac cag agt cat tgg	435
Gln Asn Asn Ala His Arg Asp Lys Glu Gly Asp Asp Gln Ser His Trp	
90 95 100	
 cgc tat gga ggc gac ccg ccc tgg ccc cgg gtg tcc cca gcc tgc gcg	483
Arg Tyr Gly Gly Asp Pro Pro Trp Pro Arg Val Ser Pro Ala Cys Ala	
105 110 115 120	
 ggc cgc ttc cag tcc ccg gtg gat atc cgc ccc cag ctc gcc gcc ttc	531
Gly Arg Phe Gln Ser Pro Val Asp Ile Arg Pro Gln Leu Ala Ala Phe	
125 130 135	
 tgc ccg gcc ctg cgc ccc ctg gaa ctc ctg ggc ttc cag ctc ccg ccg	579
Cys Pro Ala Leu Arg Pro Leu Glu Leu Leu Gly Phe Gln Leu Pro Pro	
140 145 150	
 ctc cca gaa ctg cgc ctg cgc aac aat ggc cac agt gtg caa ctg acc	627
Leu Pro Glu Leu Arg Leu Arg Asn Asn Gly His Ser Val Gln Leu Thr	
155 160 165	
 ctg cct cct ggg cta gag atg gct ctg ggt ccc ggg cgg gag tac cgg	675
Leu Pro Pro Gly Leu Glu Met Ala Leu Gly Pro Gly Arg Glu Tyr Arg	
170 175 180	
 gct ctg cag ctg cat ctg cac tgg ggg gct gca ggt cgt ccg ggc tcg	723
Ala Leu Gln Leu His Leu His Trp Gly Ala Ala Gly Arg Pro Gly Ser	
185 190 195 200	

gag cac act gtg gaa ggc cac cgt ttc cct gcc gag atc cac gtg gtt	771
Glu His Thr Val Glu Gly His Arg Phe Pro Ala Glu Ile His Val Val	
205 210 215	
cac ctc agc acc gcc ttt gcc aga gtt gac gag gcc ttg ggg cgc ccg	819
His Leu Ser Thr Ala Phe Ala Arg Val Asp Glu Ala Leu Gly Arg Pro	
220 225 230	
gga ggc ctg gcc gtg ttg gcc gcc ttt ctg gag gag ggc ccg gaa gaa	867
Gly Gly Leu Ala Val Leu Ala Ala Phe Leu Glu Glu Gly Pro Glu Glu	
235 240 245	
aac agt gcc tat gag cag ttg ctg tct cgc ttg gaa gaa atc gct gag	915
Asn Ser Ala Tyr Glu Gln Leu Leu Ser Arg Leu Glu Glu Ile Ala Glu	
250 255 260	
gaa ggc tca gag act cag gtc cca gga ctg gac ata tct gca ctc ctg	963
Glu Gly Ser Glu Thr Gln Val Pro Gly Leu Asp Ile Ser Ala Leu Leu	
265 270 275 280	
ccc tct gac ttc agc cgc tac ttc caa tat gag ggg tct ctg act aca	1011
Pro Ser Asp Phe Ser Arg Tyr Phe Gln Tyr Glu Gly Ser Leu Thr Thr	
285 290 295	
ccg ccc tgt gcc cag ggt gtc atc tgg act gtg ttt aac cag aca gtg	1059
Pro Pro Cys Ala Gln Gly Val Ile Trp Thr Val Phe Asn Gln Thr Val	
300 305 310	
atg ctg agt gct aag cag ctc cac acc ctc tct gac acc ctg tgg gga	1107
Met Leu Ser Ala Lys Gln Leu His Thr Leu Ser Asp Thr Leu Trp Gly	
315 320 325	
cct ggt gac tct cgg cta cag ctg aac ttc cga gcg acg cag cct ttg	1155
Pro Gly Asp Ser Arg Leu Gln Leu Asn Phe Arg Ala Thr Gln Pro Leu	
330 335 340	
aat ggg cga gtg att gag gcc tcc ttc cct gct gga gtg gac agc agt	1203
Asn Gly Arg Val Ile Glu Ala Ser Phe Pro Ala Gly Val Asp Ser Ser	
345 350 355 360	
cct cgg gct gct gag cca gtc cag ctg aat tcc tgc ctg gct gct ggt	1251
Pro Arg Ala Ala Glu Pro Val Gln Leu Asn Ser Cys Leu Ala Ala Gly	
365 370 375	
gac atc cta gcc ctg gtt ttt ggc ctc ctt ttt gct gtc acc agc gtc	1299
Asp Ile Leu Ala Leu Val Phe Gly Leu Leu Phe Ala Val Thr Ser Val	
380 385 390	

gcg ttc ctt gtg cag atg aga agg cag cac aga agg gga acc aaa ggg 1347
 Ala Phe Leu Val Gln Met Arg Arg Gln His Arg Arg Gly Thr Lys Gly
 395 400 405

ggg gtg agc tac cgc cca gca gag gta gcc gag act gga gcc 1389
 Gly Val Ser Tyr Arg Pro Ala Glu Val Ala Glu Thr Gly Ala
 410 415 420

tagaggctgg atcttgagaga atgtgagaag ccagccagag gcattctgagg gggagccggg 1449

aactgtcctg tctgtctcat tatgccactt ccttttaact gccaaagaaat tttttaaaat 1509

aaatatttat aat 1522

<210> 2
 <211> 459
 <212> PRT
 <213> HUMAN

<400> 2
 Met Ala Pro Leu Cys Pro Ser Pro Trp Leu Pro Leu Leu Ile Pro Ala
 -35 -30 -25

Pro Ala Pro Gly Leu Thr Val Gln Leu Leu Leu Ser Leu Leu Leu Leu
 -20 -15 -10

Met Pro Val His Pro Gln Arg Leu Pro Arg Met Gln Glu Asp Ser Pro
 -5 -1 1 5 10

Leu Gly Gly Gly Ser Ser Gly Glu Asp Asp Pro Leu Gly Glu Glu Asp
 15 20 25

Leu Pro Ser Glu Glu Asp Ser Pro Arg Glu Glu Asp Pro Pro Gly Glu
 30 35 40

Glu Asp Leu Pro Gly Glu Glu Asp Leu Pro Gly Glu Glu Asp Leu Pro
 45 50 55

Glu Val Lys Pro Lys Ser Glu Glu Glu Gly Ser Leu Lys Leu Glu Asp
 60 65 70 75

Leu Pro Thr Val Glu Ala Pro Gly Asp Pro Gln Glu Pro Gln Asn Asn
 80 85 90

Ala His Arg Asp Lys Glu Gly Asp Asp Gln Ser His Trp Arg Tyr Gly
 95 100 105

Gly Asp Pro Pro Trp Pro Arg Val Ser Pro Ala Cys Ala Gly Arg Phe
 110 115 120

Gln Ser Pro Val Asp Ile Arg Pro Gln Leu Ala Ala Phe Cys Pro Ala
 125 130 135

Leu Arg Pro Leu Glu Leu Leu Gly Phe Gln Leu Pro Pro Leu Pro Glu
 140 145 150 155

Leu Arg Leu Arg Asn Asn Gly His Ser Val Gln Leu Thr Leu Pro Pro
 160 165 170

Gly Leu Glu Met Ala Leu Gly Pro Gly Arg Glu Tyr Arg Ala Leu Gln
 175 180 185

Leu His Leu His Trp Gly Ala Ala Gly Arg Pro Gly Ser Glu His Thr
 190 195 200

Val Glu Gly His Arg Phe Pro Ala Glu Ile His Val Val His Leu Ser
 205 210 215

Thr Ala Phe Ala Arg Val Asp Glu Ala Leu Gly Arg Pro Gly Gly Leu
 220 225 230 235

Ala Val Leu Ala Ala Phe Leu Glu Glu Gly Pro Glu Glu Asn Ser Ala
 240 245 250

Tyr Glu Gln Leu Leu Ser Arg Leu Glu Glu Ile Ala Glu Glu Gly Ser
 255 260 265

Glu Thr Gln Val Pro Gly Leu Asp Ile Ser Ala Leu Leu Pro Ser Asp
 270 275 280

Phe Ser Arg Tyr Phe Gln Tyr Glu Gly Ser Leu Thr Thr Pro Pro Cys
 285 290 295

Ala Gln Gly Val Ile Trp Thr Val Phe Asn Gln Thr Val Met Leu Ser
 300 305 310 315

Ala Lys Gln Leu His Thr Leu Ser Asp Thr Leu Trp Gly Pro Gly Asp
 320 325 330

Ser Arg Leu Gln Leu Asn Phe Arg Ala Thr Gln Pro Leu Asn Gly Arg
 335 340 345

Val Ile Glu Ala Ser Phe Pro Ala Gly Val Asp Ser Ser Pro Arg Ala
 350 355 360

Ala Glu Pro Val Gln Leu Asn Ser Cys Leu Ala Ala Gly Asp Ile Leu
 365 370 375

Ala Leu Val Phe Gly Leu Leu Phe Ala Val Thr Ser Val Ala Phe Leu
 380 385 390 395

Val Gln Met Arg Arg Gln His Arg Arg Gly Thr Lys Gly Gly Val Ser
 400 405 410

Tyr Arg Pro Ala Glu Val Ala Glu Thr Gly Ala
 415 420

<210> 3

<211> 29

<212> DNA

<213> HUMAN

<400> 3

cgcccagtggt gtcattcttc ccagaagag

29

<210> 4

<211> 19

<212> DNA

<213> HUMAN

<400> 4

ggaatcctcc tgcattccgg

19

<210> 5

<211> 10898

<212> DNA

<213> HUMAN

<220>

<221> gene

<222> (1) .. (10898)

<400> 5

ggatcctgtt gactcgtgac cttaccccca accctgtgct ctctgaaaca tgagctgtgt 60

ccactcaggg ttaaattgat taagggcggt gcaagatgtg ctttggtaaa cagatgcttg 120

aaggcagcat gctcgttaag agtcattacc aatccctaatt ctcaagtaatt cagggacaca 180

aacactgcgg aaggccgcag ggctctctgc ctaggaaaac cagagacctt tgttcacttg 240
tttatctgac cttccctcca ctattgtcca tgaccctgcc aaatccccct ctgtgagaaa 300
cacccaagaa ttatcaataa aaaaataaat ttaaaaaaaaa aatacaaaaa aaaaaaaaaa 360
aaaaaaaaaa gacttacgaa tagttattga taaatgaata gctattggta aagccaagta 420
aatgatcata ttcaaaacca gacggccatc atcacagctc aagtctacct gatttgatct 480
ctttatcatt gtcattcttt ggattcacta gattagtcac catcctcaaa attctcccc 540
aagttctaata tacgttccaa acatttaggg gttacatgaa gcttgaacct actaccttct 600
ttgcttttga gccatgagtt gtaggaatga tgagtttaca ccttacatgc tggggattaa 660
tttaaaacttt acctctaagt cagttgggta gcctttggct tatttttgta gctaattttg 720
tagttaatgg atgcactgtg aatcttgcta tgatagtttt cctccacact ttgccactag 780
gggtaggtag gtactcagtt ttcagtaatt gcttacctaa gaccctaagc cctatttctc 840
ttgtactggc ctttatctgt aatatgggca tatttaatac aatataattt ttggagtttt 900
tttgtttggt tgtttgtttg tttttttgag acggagtctt gcacctgtca tgcccaggct 960
ggagtagcag tggtgccatc tcggctcact gcaagctcca cctcccgagt tcacgccatt 1020
ttcctgcctc agcctccga gtagctggga ctacaggcgc ccgccaccat gcccggttaa 1080
ttttttgtat ttttggtaga gacgggggtt caccgtgta gccagaatgg tctcgatctc 1140
ctgacttcgt gatccacccg cctcggcctc ccaaagttct gggattacag gtgtgagcca 1200
ccgcacctgg ccaatttttt gagtctttta aagtaaaaat atgtcttgta agctggtaac 1260
tatggtacat ttccttttat taatgtgggt ctgacgggtca tataggttct tttgagtttg 1320
gcatgcatat gctacttttt gcagtccttt cattacattt ttctctcttc atttgaagag 1380
catgttatat cttttagctt cacttggctt aaaaggttct ctcatagcc taacacagt 1440
tcattgttgg taccacttgg atcataagt gaaaaacagt caagaaattg cacagtaata 1500
cttgtttgta agagggatga ttcaggtgaa tctgacacta agaaactccc ctacctgagg 1560
tctgagattc ctctgacatt gctgtatata ggcttttctt ttgacagcct gtgactgcgg 1620

actatttttc ttaagcaaga tatgctaaag ttttgtgagc ctttttccag agagaggtct 1680
catatctgca tcaagtgaga acatataatg tctgcatggt tccatatttc aggaatgttt 1740
gcttgtgttt tatgctttta tatagacagg gaaacttggt cctcagtgac ccaaagagg 1800
tggaattgt tattggatat catcattggc ccacgctttc tgaccttgga aacaattaag 1860
ggttcataat ctcaattctg tcagaattgg tacaagaaat agctgctatg tttcttgaca 1920
ttccacttgg taggaaataa gaatgtgaaa ctcttcaggt ggtgtgtgtc cctngttttt 1980
ttgcaatttc cttcttactg tgttaaaaaa aagtatgatc ttgctctgag aggtgaggca 2040
ttcttaatca tgatctttaa agatcaataa tataatcctt tcaaggatta tgtctttatt 2100
ataataaaga taatttgtct ttaacagaat caataatata atcccttaa ggattatatc 2160
tttgctgggc gcagtggctc acacctgtaa tccagcact ttgggtggcc aagggtggaag 2220
gatcaaattt gcctacttct atattatctt ctaaagcaga attcatctct cttccctcaa 2280
tatgatgata ttgacagggt ttgccctcac tcactagatt gtgagctcct gctcagggca 2340
ggtagcggtt tttgtttttg tttttgtttt tcttttttga gacagggtct tgctctgtca 2400
cccaggccag agtgcaatgg tacagtctca gctcactgca gcctcaaccg cctcggtctca 2460
aaccatcatc ccatttcagc ctctgagta gctgggacta caggcacatg ccattacacc 2520
tggctaattt ttttgtattt ctagtagaga cagggtttgg ccatgttgcc cgggctggtc 2580
tcgaactcct ggactcaagc aatccacca cctcagcctc ccaaagttag ggaccgtgtc 2640
ttattcattt ccatgtccct agtccatagc ccagtgtgtg acctatggta gtactaaata 2700
aatatttggt gaatgcaata gtaaatagca tttcaggagg caagaactag attaacaaag 2760
gtggtaaaag gtttgagaa aaaaataata gtttaatttg gctagagtat gagggagagt 2820
agtaggagac aagatggaaa ggtctcttgg gcaaggtttt gaaggaagtt ggaagtcaga 2880
agtacacaat gtgcatatcg tggcaggcag tggggagcca atgaaggctt ttgagcagga 2940
gagtaatgtg ttgaaaaata aatataggtt aaacctatca gagccctct gacacataca 3000
cttgcttttc attcaagctc aagtttgtct cccacatacc cattacttaa ctcaccctcg 3060

ggctccccta gcagcctgcc ctacctcttt acctgcttcc tggaggagtc agggatgtat 3120
acatgagctg ctttccctct cagccagagg acatgggggg cccagctcc cctgcctttc 3180
cccttctgtg cctggagctg ggaagcaggc cagggttagc tgaggctggc tggcaagcag 3240
ctgggtgggtg ccaggagag cctgcatagt gccagggtgg gccttggggt ccaagctagt 3300
ccatggcccc gataaccttc tgctgtgca cacacctgcc cctcactcca cccccatcct 3360
agctttggta tgggggagag ggcacagggc cagacaaacc tgtgagactt tggctccatc 3420
tctgcaaaag ggcgctctgt gagtacgct gctccctcc aggcttgctc ctccccacc 3480
cagctctcgt ttccaatgca cgtacagccc gtacacaccg tgtgctggga caccacacag 3540
tcagccgcat ggctcccctg tgcccagcc cctggctccc tctgttgatc ccggcccctg 3600
ctccaggcct cactgtgcaa ctgctgctgt cactgctgct tctgggtgct gtccatcccc 3660
agaggttgcc ccggatgcag gaggattccc ccttgggagg aggctcttct ggggaagatg 3720
accactggg cgaggaggat ctgcccagtg aagaggattc accagagag gaggatccac 3780
ccggagagga ggatctacct ggagaggagg atctacctgg agaggaggat ctacctgaag 3840
ttaagcctaa atcagaagaa gagggtccc tgaagttaga ggatctacct actgttgagg 3900
ctcctggaga tcctcaagaa cccagaata atgccacag ggacaaagaa ggtaagtgg 3960
catcaatctc caaatccagg ttccaggagg ttcatgactc ccctcccata cccagccta 4020
ggctctgttc actcaggga ggaggggaga ctgtactccc cacagaagcc cttccagagg 4080
tcccatacca atatcccat cccactctc ggaggtagaa agggacagat gtggagagaa 4140
aataaaaagg gtgcaaaagg agagaggtaga gctggatgag atgggagaga agggggaggc 4200
tggaagagag aaaggatga gaactgcaga tgagagaaaa aatgtgcaga cagaggaaaa 4260
aaataggtgg agaaggagag tcagagagtt tgaggggaag agaaaaggaa agcttgggag 4320
gtgaagtggg taccagagac aagcaagaag agctggtaga agtcatctca tcttaggcta 4380
caatgaggaa ttgagaccta ggaagaaggg acacagcagg tagagaaacg tggcttcttg 4440
actcccaagc caggaatttg gggaaagggg ttggagacca tacaaggcag agggatgagt 4500

ggggagaaga aagaaggag aaaggaaaga tgggtgtactc actcatttgg gactcaggac 4560
tgaagtgcc actcactttt tttttttttt tttttgagac aaactttcac ttttgttgcc 4620
caggctggag tgcaatggcg cgatctcggc tcaactgcaac ctccacctcc cgggttcaag 4680
tgattctcct gcctcagcct ctagccaagt agctgcgatt acaggcatgc gccaccacgc 4740
ccggctaatt tttgtatttt tagtagagac ggggtttcgc catgttggtc aggctgggtc 4800
cgaactcctg atctcagggt atccaaccac cctggcctcc caaagtgctg ggattatagg 4860
cgtgagccac agcgccctggc ctgaagcagc cactcacttt tacagaccct aagacaatga 4920
ttgcaagctg gtaggattgc tgtttggccc acccagctgc ggtgttgagt ttgggtgcgg 4980
tctcctgtgc tttgcacctg gcccgcttaa ggcatttgtt acccgtaatg ctctgtgaag 5040
gcatctgcgt ttgtgacatc gttttggtcg ccaggaaggg attggggctc taagcttgag 5100
cggttcatcc ttttcattta tacaggggat gaccagagtc attggcgcta tggagggtgag 5160
acaccacccc gctgcacaga cccaatctgg gaaccagct ctgtggatct cccctacagc 5220
cgtccctgaa cactgggtccc gggcgctcca cccgcgcgcc accgtcccac cccctcacct 5280
tttctacccg gggtccctaa gttcctgacc taggcgtcag acttcctcac tatactctcc 5340
caccacaggc gaccgcctt ggccccgggt gtccccagcc tgcgcgggcc gcttccagtc 5400
cccggtggat atccgcccc agctcgccgc cttctgcccc gccctgcgcc ccctggaact 5460
cctgggcttc cagctcccc cgctcccaga actgcgcctg cgcaacaatg gccacagtgg 5520
tgagggggtc tccccgccga gacttgggga tggggcgggg cgcagggaag ggaaccgtcg 5580
cgcagtgcct gcccgggggg tgggctggcc ctaccgggcg gggccggctc acttgccctc 5640
ccctacgcag tgcaactgac cctgcctcct gggctagaga tggctctggg tcccgggcgg 5700
gagtaccggg ctctgcagct gcatctgcac tggggggctg caggctcgtc gggctcggag 5760
cacactgtgg aaggccaccg tttccctgcc gaggtgagcg cggactggcc gagaaggggc 5820
aaaggagcgg ggcggacggg ggccagagac gtggccctct cctaccctcg tgtccttttc 5880
agatccacgt ggttcacctc agcaccgcct ttgccagagt tgacgaggcc ttggggcgcc 5940

cgaggaggcct ggccgtgttg gccgcctttc tggagggtacc agatcctgga caccocctac 6000
tccccgcttt cccatcccat gtcctccccg gactctatcg tggagccaga gaccccatcc 6060
cagcaagctc actcaggccc ctggctgaca aactcattca cgcactgttt gttcatttaa 6120
caccactgt gaaccaggca ccagccccc acaaggattc tgaagctgta ggtccttgcc 6180
tctaaggagc ccacagccag tgggggaggc tgacatgaca gacacatagg aaggacatag 6240
taaagatggt ggtcacagag gaggtgacac ttaaagcctt cactggtaga aaagaaaagg 6300
aggtgttcat tgcagaggaa acagaatgtg caaagactca gaatatggcc tatttaggga 6360
atggctacat acaccatgat tagaggaggc ccagtaaagg gaagggatgg tgagatgcct 6420
gctaggttca ctcaactcact tttatttatt tatttatttt ttgacagtc tctctgtcgc 6480
ccaggctgga gtgcagtggg gtgatcttgg gtcactgcaa cttccgcctc ccgggttcaa 6540
gggattctcc tgcctcagct tcctgagtag ctgggggttac aggtgtgtgc caccatgcc 6600
agctaatttt tttttgtatt tttagtagac agggtttcac catgttgggc aggtgtgtct 6660
caaactcctg gcctcaagtg atccgcctga ctcaagcctac caaagtgtg attacaagt 6720
tgagccaccg tgcccagcca cactcactga ttctttaatg ccagccacac agcacaaagt 6780
tcagagaaat gcctccatca tagcatgtca atatgttcat actcttaggt tcatgatgtt 6840
cttaacatta ggttcataag caaaataaga aaaaagaata ataaataaaa gaagtggcat 6900
gtcaggacct cacctgaaaa gccaaacaca gaatcatgaa ggtgaatgca gaggtgacac 6960
caacacaaag gtgtatatat ggtttctgt ggggagtatg tacggaggca gcagtgagt 7020
agactgcaaa cgtcagaagg gcacgggtca ctgagagcct agtatcctag taaagtgggc 7080
tctctccctc tctctccagc ttgtcattga aaaccagtcc accaagcttg ttggttcgca 7140
cagcaagagt acatagagtt tgaaataata cataggattt taagaggag acactgtctc 7200
taaaaaaaaa aacaacagca acaacaaaaa gcaacaacca ttacaatttt atgttccctc 7260
agcattctca gagctgagga atgggagagg actatgggaa ccccttcat gttccggcct 7320
tcagccatgg ccctggatac atgcactcat ctgtcttaca atgtcattcc ccaggagg 7380

cccggaagaa aacagtgcct atgagcagtt gctgtctcgc ttggaagaaa tcgctgagga 7440
aggtcagttt gttgggtctgg ccactaatct ctgtggccta gttcataaag aatcaccctt 7500
tgagagcttca ggtctgaggc tggagatggg ctccctccag tgcaggaggg attgaagcat 7560
gagccagcgc tcattcttgat aataaccatg aagctgacag acacagttac ccgcaaacgg 7620
ctgcctacag attgaaaacc aagcaaaaac cgccgggcac ggtggctcac gcctgtaatc 7680
ccagcacttt gggaggccaa ggcaggtgga tcacgaggtc aagagatcaa gaccatcctg 7740
gccaacatgg tgaaacccca tctctactaa aaatacga aaatagccag gcgtgggtggc 7800
gggtgcctgt aatcccagct actcgggagg ctgaggcagg agaatggcat gaaccgggga 7860
ggcagaagtt gcagtgagcc gagatcgtgc cactgcactc cagcctgggc aacagagcga 7920
gactcttgtc tcaaaaaaaaa aaaaaaaaaa gaaaaccaag caaaaaccaa aatgagacaa 7980
aaaaaacaag accaaaaaat ggtgttttga aattgtcaag gtcaagtctg gagagctaaa 8040
ctttttctga gaactgttta tctttaataa gcatcaata ttttaacttt gtaaatactt 8100
ttgttggaat tcgttctctt cttagtcact cttgggtcat tttaaatctc acttactcta 8160
ctagaccttt taggtttctg ctagactagg tagaactctg cttttgcatt tcttgtgtct 8220
gttttgata gttatcaata ttcatttta tttacaagtt attcagatca ttttttcttt 8280
tctttttttt tttttttttt ttttttacat ctttagtaga gacagggttt caccatattg 8340
gccaggctgc tctcaaactc ctgacctgt gatccaccag cctcggcctc ccaaagtgtc 8400
gggattcatt ttttcttttt aatttgcctt gggcttaaac ttgtggccca gcactttatg 8460
atggtacaca gagttaagag tgtagactca gacggctctt cttctttcct tctcttcctt 8520
cctcccttcc ctcccacctt cctttctctc cttcctttct ttcttctctt cttgcttctt 8580
caggcctctt ccagttgctc caaagccctg tacttttttt tgagttaacg tcttatggga 8640
agggcctgca cttagtgaag aagtggctct agagttgagt taccttggct tctgggaggt 8700
gaaactgtat ccctataccc tgaagcttta agggggtgca atgtagatga gacccaaca 8760
tagatcctct tcacaggctc agagactcag gtcccaggac tggacatctc tgcactcctg 8820

ccctctgact tcagccgcta cttccaatat gaggggtctc tgactacacc gccctgtgcc 8880
caggggtgtca tctggactgt gtttaaccag acagtgatgc tgagtgctaa gcaggtgggc 8940
ctgggggtgtg tgtggacaca gtgggtgcgg gggaaagagg atgtaagatg agatgagaaa 9000
caggagaaga aagaaatcaa ggctgggctc tgtggcttac gcctataatc ccaccacgtt 9060
gggaggctga ggtgggagaa tggtttgagc ccaggagttc aagacaaggc ggggcaacat 9120
agtgtgacct catctctacc aaaaaaaccc caacaaaacc aaaaatagcc gggcatgggtg 9180
gtatgcggcc tagtcccagc tactcaagga ggctgaggtg ggaagatcgc ttgattccag 9240
gagtttgaga ctgcagtgag ctatgatccc accactgcct accatcttta ggatacattt 9300
atttatttat aaaagaaatc aagaggctgg atggggaata caggagctgg aggggtggagc 9360
cctgaggtgc tggttgtgag ctggcctggg acccttgttt cctgtcatgc catgaacca 9420
cccacactgt cactgacct ccctagctcc acaccctctc tgacacctg tggggacctg 9480
gtgactctcg gctacagctg aacttccgag cgacgcagcc tttgaatggg cgagtgattg 9540
aggcctcctt ccctgctgga gtggacagca gtccctgggc tgctgagcca ggtacagctt 9600
tgtctggttt cccccagcc agtagtcctt taccctccca tgtgtgtgcc agtgtctgtc 9660
attggtggtc acagcccgcc tctcacatct cctttttctc tccagtccag ctgaattcct 9720
gcctggctgc tggtgagtct gccctcctc ttggctctga tgccaggaga ctctcagca 9780
ccattcagcc ccagggtgc tcaggaccgc ctctgctccc tctccttttc tgcagaacag 9840
acccaacccc caatattaga gaggcagatc atgggtgggga ttccccatt gtccccagag 9900
gctaattgat tagaatgaag cttgagaaat ctcccagcat ccctctcgca aaagaatccc 9960
ccccctttt tttaaagata gggctctact ctgtttgccc caggctgggg tgttgtggca 10020
cgatcatagc tcactgcagc ctcgaaactc taggctcagg caatcctttc accttagctt 10080
ctcaaagcac tgggactgta ggcatgagcc actgtgcctg gccccaaacg gcccttttac 10140
ttggctttta ggaagcaaaa acggtgctta tcttaccct tctcgtgtat ccaccctcat 10200
cccttggtg gcctcttctg gagactgagg cactatgggg ctgcctgaga actcggggca 10260

ggggtggtgg agtgcactga ggcaggtgtt gaggaactct gcagaccctt cttccttccc 10320
 aaagcagccc tctctgctct ccacgcagg tgacatccta gccctggttt ttggcctcct 10380
 ttttgcgtgc accagcgctg cgttccttgt gcagatgaga aggcagcaca ggtattacac 10440
 tgaccctttc ttcaggcaca agcttcccc acccttgttg agtcacttca tgcaaagcgc 10500
 atgcaaata gctgctcctg ggccagtttt ctgattagcc tttcctgttg tgtacacaca 10560
 gaaggggaac caaagggggg gtgagctacc gccagcaga ggtagccgag actggagcct 10620
 agaggctgga tcttgagaa tgtgagaagc cagccagagg catctgaggg ggagccggta 10680
 actgtcctgt cctgctcatt atgccacttc cttttaactg ccaagaaatt ttttaaaata 10740
 aatatttata ataaaatatg tgtagtcac ctttggtccc caaatcagaa ggaggtattt 10800
 gaatttccta ttactgttat tagcaccaat ttagtggtaa tgcatttatt ctattacagt 10860
 tcggcctcct tccacacatc actccaatgt gttgctcc 10898

<210> 6

<211> 37

<212> PRT

<213> HUMAN

<400> 6

Met Ala Pro Leu Cys Pro Ser Pro Trp Leu Pro Leu Leu Ile Pro Ala

1

5

10

15

Pro Ala Pro Gly Leu Thr Val Gln Leu Leu Leu Ser Leu Leu Leu

20

25

30

Met Pro Val His Pro

35

<210> 7

<211> 25

<212> DNA

<213> HUMAN

<400> 7

tggggttctt gaggatctcc aggag

25

<210> 8

<211> 26

<212> DNA

<213> HUMAN

<400> 8

ctctaacttc agggagccct cttctt

26

<210> 9

<211> 48

<212> DNA

<213> HUMAN

<220>

<221> primer_bind

<222> (1)..(48)

<400> 9

cuacuacuac uaggccacgc gtcgactagt acgggnnggg nngggng

48

<210> 10

<211> 6

<212> PRT

<213> HUMAN

<400> 10

Glu Glu Asp Leu Pro Ser

1

5

<210> 11

<211> 6

<212> PRT

<213> HUMAN

<400> 11

Gly Glu Asp Asp Pro Leu

1

5

<210> 12

<211> 21

<212> PRT

<213> HUMAN

<400> 12

Asn Asn Ala His Arg Asp Lys Glu Gly Asp Asp Gln Ser His Trp Arg
 1 5 10 15

Tyr Gly Gly Asp Pro
 20

<210> 13

<211> 16

<212> PRT

<213> HUMAN

<400> 13

His Pro Gln Arg Leu Pro Arg Met Gln Glu Asp Ser Pro Leu Gly Gly
 1 5 10 15

<210> 14

<211> 24

<212> PRT

<213> HUMAN

<400> 14

Glu Glu Asp Ser Pro Arg Glu Glu Asp Pro Pro Gly Glu Glu Asp Leu
 1 5 10 15

Pro Gly Glu Glu Asp Leu Pro Gly
 20

<210> 15

<211> 13

<212> PRT

<213> HUMAN

<400> 15

Leu Glu Glu Gly Pro Glu Glu Asn Ser Ala Tyr Glu Gln
 1 5 10

<210> 16

<211> 16

<212> PRT

<213> HUMAN

<400> 16

Met Arg Arg Gln His Arg Arg Gly Thr Lys Gly Gly Val Ser Tyr Arg
 1 5 10 15

<210> 17
<211> 45
<212> DNA
<213> HUMAN

<400> 17
gtcgttagct ccatgggtca tatgcagagg ttgccccgga tgcag 45

<210> 18
<211> 43
<212> DNA
<213> HUMAN

<400> 18
gaagatctct tactcgagca ttctccaaga tccagcctct agg 43

<210> 19
<211> 10
<212> DNA
<213> HUMAN

<400> 19
ctccatctct 10

<210> 20
<211> 10
<212> DNA
<213> HUMAN

<400> 20
ccacccccat 10

<210> 21
<211> 205
<212> DNA
<213> HUMAN

<400> 21
acctgccccct cactccaccc ccatcctagc tttggtatgg gggagagggc acagggccag 60
acaaacctgt gagacttttg ctccatctct gcaaaagggc gctctgtgag tcagcctgct 120

ccccctccagg cttgtctctc ccccacccag ctctcgtttc caatgcacgt acagcccgta 180

cacaccgtgt gctgggacac cccac

205

<210> 22

<211> 8

<212> PRT

<213> HUMAN

<400> 22

Leu Glu His His His His His His

1

5

<210> 23

<211> 10

<212> DNA

<213> HUMAN

<220>

<221> misc_feature

<222> (1)..(10)

<400> 23

yyycayyyyy

10

<210> 24

<211> 10

<212> DNA

<213> HUMAN

<300>

<301> Locker and Buzard,

<303> DNA Sequencing and Mapping

<304> 1

<306> 3-11

<307> 1990

<400> 24

tgtgagactt

10

<210> 25

<211> 4

<212> PRT

<213> HUMAN

<220>

<221> SITE

<222> (1) .. (4)

<400> 25

Ser Pro Xaa Xaa

1

<210> 26

<211> 4

<212> PRT

<213> HUMAN

<220>

<221> SITE

<222> (1) .. (4)

<400> 26

Thr Pro Xaa Xaa

1

<210> 27

<211> 540

<212> DNA

<213> HUMAN

<220>

<221> promoter

<222> (1) .. (540)

<400> 27

cttgccttttc attcaagctc aagtttgtct cccacatacc cattacttaa ctcaccctcg 60

ggctccccta gcagcctgcc ctacctcttt acctgcttcc tgggtggagtc agggatgtat 120

acatgagctg ctttccctct cagccagagg acatgggggg cccagctcc cctgcctttc 180

cccttctgtg cctggagctg ggaagcaggc caggggttagc tgaggctggc tggcaagcag 240

ctgggtggtg ccaggagag cctgcatagt gccagggtggt gccttggggtt ccaagctagt 300

ccatggcccc gataaccttc tgctgtgca cacacctgcc cctcactcca cccccatcct 360

agcttttgta tgggggagag ggcacagggc cagacaaacc tgtgagactt tggctccatc 420

tctgcaaaag ggcgctctgt gagtcagcct gctccccctcc aggcttgctc ctccccacc 480

cagctctcgt ttccaatgca cgtacagccc gtacacaccg tgtgctggga caccacacag 540

<210> 28

<211> 445

<212> DNA

<213> HUMAN

<220>

<221> exon

<222> (1)

<223> 1st MN exon

<400> 28

gcccgtagac accgtgtgct gggacacccc acagtcagcc gcatggctcc 50

cctgtgcccc agccccctggc tccctctgtt gatccccggcc cctgtctcag 100

gcctcactgt gcaactgctg ctgtcactgc tgcttctggg gcctgtccat 150

ccccagagggt tgccccggat gcaggaggat tcccccttgg gaggaggctc 200

ttctggggaa gatgaccac tgggcgagga ggatctgccc agtgaagagg 250

attcaccacag agaggaggat ccacccggag aggaggatct acctggagag 300

gaggatctac ctggagagga ggatctacct gaagttaagc ctaaatacaga 350

agaagagggc tccctgaagt tagaggatct acctactgtt gaggctcctg 400

gagatcctca agaaccacag aataatgccc acagggacaa agaag 445

<210> 29

<211> 30

<212> DNA

<213> HUMAN

<220>

<221> exon

<222> (1)

<223> 2nd MN exon

<400> 29

gggatgacca gagtcattgg cgctatggag 30

<210> 30
 <211> 171
 <212> DNA
 <213> HUMAN

<220>
 <221> exon
 <222> (1)
 <223> 3rd MN exon

<400> 30	
gcgaccgcgc ctggccccgg gtgtccccag cctgcgcggg ccgcttccag	50
tccccgggtgg atatccgccc ccagctcgcc gccttctgcc cggccctgcg	100
ccccctggaa ctctctgggct tccagctccc gccgctccca gaactgcgcc	150
tgcgcaacaa tggccacagt g	171

<210> 31
 <211> 143
 <212> DNA
 <213> HUMAN

<220>
 <221> exon
 <222> (1)
 <223> 4th MN exon

<400> 31	
tgcaactgac cctgcctcct gggctagaga tggctctygg tcccggggcgg	50
gagtaccggg ctctgcagct gcattctgcac tgggggggctg caggctcgtec	100
gggctcggag cacactgtgg aaggccaccg tttccctgcc gag	143

<210> 32
 <211> 93
 <212> DNA
 <213> HUMAN

<220>
 <221> exon
 <222> (1)
 <223> 5th MN exon

<400> 32

atccacgtgg ttcacctcag caccgccttt gccagagttg acgaggcctt 50

ggggcgcccc ggaggcctgg ccgtgttggc cgcctttctg gag 93

<210> 33

<211> 67

<212> DNA

<213> HUMAN

<220>

<221> exon

<222> (1)

<223> 6th MN exon

<400> 33

gagggccccg aagaaaacag tgcctatgag cagttgctgt ctgccttgga 50

agaaatcgct gaggaag 67

<210> 34

<211> 158

<212> DNA

<213> HUMAN

<220>

<221> exon

<222> (1)

<223> 7th MN exon

<400> 34

gtcagagac tcaggtecca ggactggaca tatctgcaact cctgccctct 50

gacttcagcc gctacttcca atatgagggg tctctgacta caccgcctg 100

tgcccagggt gtcacttgga ctgtgtttta ccagacagtg atgctgagt ctaagcag 158

<210> 35

<211> 145

<212> DNA

<213> HUMAN

<220>

<221> exon

<222> (1)

<223> 8th MN exon

<400> 35

ctccacaccc tctctgacac cctgtgggga cctggtgact ctcggtaca 50

gctgaacttc cgagcgacgc agcctttgaa tgggcgagtg attgaggcct 100

ccttcctgc tggagtggac agcagtcctc gggctgctga gccag 145

<210> 36

<211> 27

<212> DNA

<213> HUMAN

<220>

<221> exon

<222> (1)

<223> 9th MN exon

<400> 36

tccagctgaa ttctgcctg gctgctg 27

<210> 37

<211> 82

<212> DNA

<213> HUMAN

<220>

<221> exon

<222> (1)

<223> 10th MN exon

<400> 37

gtgacatcct agccctgggt tttggcctcc tttttgctgt caccagcgtc 50

gcgttccttg tgcagatgag aaggcagcac ag 82

<210> 38

<211> 191

<212> DNA

<213> HUMAN

<220>

<221> exon

<222> (1)

<223> 11th MN exon

<400> 38

```

aaggggaacc aaaggggtg tgagctaccg cccagcagag gtagccgaga      50
ctggagccta gaggctggat cttggagaat gtgagaagcc agccagaggc      100
atctgagggg gagccggtaa ctgtcctgtc ctgtcatta tgccacttcc      150
ttttaactgc caagaaattt tttaaaataa atatttataa t                191

```

<210> 39

<211> 1174

<212> DNA

<213> HUMAN

<220>

<221> intron

<222> (1) .. (1174)

<223> 1st MN intron

<400> 39

```

gtaagtggtc atcaatctcc aaatccaggt tccaggaggt tcatgactcc cctcccatac 60
cccagcctag gctctgttca ctcaggggaag gaggggagac tgtactcccc acagaagccc 120
ttccagaggt cccataccaa tatccccatc cccactctcg gaggtagaaa gggacagatg 180
tgagagagaaa ataaaaaggg tgcaaaaagga gagaggtgag ctggatgaga tgggagagaa 240
gggggagggt ggagaagaga aagggatgag aactgcagat gagagaaaaa atgtgcagac 300
agaggaaaaa aataggtgga gaaggagagt cagagagttt gaggggaaga gaaaaggaaa 360
gcttgggagg tgaagtgggt accagagaca agcaagaaga gctggtagaa gtcactcat 420
cttaggctac aatgaggaat tgagacctag gaagaaggga cacagcaggt agagaaacgt 480
ggcttcttga ctccaagcc aggaatttgg ggaaaggggt tggagaccat acaaggcaga 540
gggatgagtg gggagaagaa agaagggaga aaggaaagat ggtgtactca ctcatattgg 600
actcaggact gaagtgccca ctcaactttt tttttttttt ttttgagaca aactttcact 660
ttgtttgccc aggctggagt gcaatggcgc gatctcgggt cactgcaacc tccacctccc 720

```

gggttcaagt gattctcctg cctcagcctc tagccaagta gctgcgatta caggcatgcg 780
ccaccacgcc eggctaattt ttgtatTTTT agtagagacg gggtttcgcc atgttggtca 840
ggctgggtctc gaactcctga tctcaggtga tccaaccacc ctggcctccc aaagtgctgg 900
gattataggc gtgagccaca gcgcctggcc tgaagcagcc actcactttt acagacccta 960
agacaatgat tgcaagctgg taggattgct gtttggccca cccagctgcg gtgttgagtt 1020
tggttgcggt ctctgtgct ttgcacctgg cccgcttaag gcatttgta cccgtaatgc 1080
tctgtgaagg catctgcgtt tgtgacatcg ttttggtcgc caggaaggga ttggggctct 1140
aagcttgagc ggttcctcct tttcatttat acag 1174

<210> 40

<211> 193

<212> DNA

<213> HUMAN

<220>

<221> intron

<222> (1)..(193)

<223> 2nd MN intron

<400> 40

gtgagacacc caccgctgc acagacccaa tctgggaacc cagctctgtg gatctccct 60
acagccgtcc ctgaacactg gtcccgggcy tcccaccgc cgcccaccgt cccacccct 120
caccttttct acccgggttc cctaagttcc tgacctaggc gtcagacttc ctactatac 180
tctcccaccc cag 193

<210> 41

<211> 131

<212> DNA

<213> HUMAN

<220>

<221> intron

<222> (1)..(131)

<223> 3rd MN intron

<400> 41

gtgaggggggt ctccccgccg agacttgggg atggggcggg gcgcagggaa gggaaccgtc 60
 gcgcagtgcc tgccccggggg ttgggctggc cctaccgggc ggggccggct cacttgctc 120
 tccctacgca g 131

<210> 42
 <211> 89
 <212> DNA
 <213> HUMAN

<220>
 <221> intron
 <222> (1)..(89)
 <223> 4th MN intron

<400> 42
 gtgagcgcgg actggccgag aaggggcaaa ggagcggggc ggacgggggc cagagacgtg 60
 gccctctcct accctcgtgt ccttttcag 89

<210> 43
 <211> 1400
 <212> DNA
 <213> HUMAN

<220>
 <221> intron
 <222> (1)..(1400)
 <223> 5th MN intron

<400> 43
 gtaccagatc ctggacaccc cctactcccc gctttcccat cccatgctcc tcccggactc 60
 tatcgtggag ccagagaccc catcccagca agctcactca ggcccctggc tgacaaactc 120
 attcacgcac tgtttgttca tttaacaccc actgtgacc aggcaccagc cccaacaag 180
 gattctgaag ctgtaggtcc ttgcctctaa ggagcccaca gccagtgggg gaggctgaca 240
 tgacagacac ataggaagga catagtaaag atggtggtca cagaggaggt gacacttaaa 300
 gccttcactg gtagaaaaga aaaggaggtg ttcattgcag aggaaacaga atgtgcaaag 360
 actcagaata tggcctatct agggaaatggc tacatacacc atgattagag gaggcccagt 420

aaaggaagg gatggtgaga tgccctgctag gttcactcac tcacttttat ttattttattt 480
atttttttga cagtctctct gtcgccagg ctggagtga gtggtgtgat cttgggtcac 540
tgcaacttcc gcctcccggg ttcaagggat tctcctgcct cagcttcctg agtagctggg 600
gttacaggtg tgtgccacca tgcccagcta attttttttt gtatttttag tagacagggg 660
ttcaccatgt tggtcaggct ggtctcaaac tcctggcctc aagtgatccg cctgactcag 720
cctaccaaag tgctgattac aagtgtgagc caccgtgcc agccacactc actgattctt 780
taatgccagc cacacagcac aaagttcaga gaaatgcctc catcatagca tgtcaatatg 840
ttcatactct taggttcatg atgttcttaa cattaggttc ataagcaaaa taagaaaaaa 900
gaataataaa taaaagaagt ggcattgtcag gacctcacct gaaaagccaa acacagaatc 960
atgaagggtga atgcagaggt gacaccaaca caaagggtga tatatgggtt cctgtgggga 1020
gtatgtacgg aggcagcagt gagtgaact gcaaactca gaagggcacg ggtcactgag 1080
agcctagtat cctagtaaag tgggctctct ccctctctct ccagcttgct attgaaaacc 1140
agtcaccaaa gcttggttgg tgcacagca agagtacata gagtttgaaa taatacatag 1200
gattttaaga gggagacact gtctctaaaa aaaaaaaca cagcaacaac aaaaagcaac 1260
aaccattaca attttatgtt ccctcagcat tctcagagct gaggaatggg agaggactat 1320
gggaaccccc ttcatgttcc ggccttcagc catggccctg gatacatgca ctcatctgtc 1380
ttacaatgtc attccccag 1400

<210> 44

<211> 1334

<212> DNA

<213> HUMAN

<220>

<221> intron

<222> (1)..(1334)

<223> 6th MN intron

<400> 44

gtcagtttgt tggctctggc actaatctct gtggcctagt tcataaagaa tcaccctttg 60

gagcttcagg tctgaggctg gagatgggct ccctccagtg caggagggat tgaagcatga 120
gccagcgctc atcttgataa taaccatgaa gctgacagac acagttaccc gcaaaccggct 180
gcctacagat tgaaaaccaa gcaaaaaccg ccgggcacgg tggctcacgc ctgtaatccc 240
agcacttttg gaggccaagg caggtggatc acgaggtcaa gagatcaaga ccatcctggc 300
caacatggtg aaaccccatc tctactaaaa atacgaaaaa atagccaggc gtggtggcgg 360
gtgcctgtaa tcccagctac tcgggaggct gaggcaggag aatggcatga acccgggagg 420
cagaagttgc agtgagccga gatcgtgcc ctgcactcca gcctgggcaa cagagcgaga 480
ctcttgctc aaaaaaaaaa aaaaaaaga aaaccaagca aaaacaaaa tgagacaaaa 540
aaaacaagac caaaaaatgg tgtttgaaa ttgtcaagg caagtctgga gagctaaact 600
tttctgaga actgtttatc ttttaataagc atcaaataatt ttaactttgt aaatactttt 660
gttggaatc gttctcttct tagtcactct tgggtcattt taaatctcac ttactctact 720
agacctttta ggtttctgct agactaggta gaactctgcc tttgcatttc ttgtgtctgt 780
tttgatatag tatcaatatt catatttatt tacaagttat tcagatcatt ttttcttttc 840
tttttttttt tttttttttt ttttacatct ttagtagaga cagggtttca ccatattggc 900
caggctgctc tcaaactcct gaccttgta tccaccagcc tcggcctccc aaagtgctgg 960
gattcatttt ttctttttta tttgctctgg gcttaaactt gtggcccagc actttatgat 1020
ggtacacaga gttaagagt tagactcaga cggtcttct tctttccttc tcttccttc 1080
tcccttcctt cccaccttc cttctctctt tctttctttt ctctctctct tgcttcctca 1140
ggcctcttc agttgctcca aagccctgta cttttttttg agttaacgct ttatgggaag 1200
ggcctgcact tagtgaagaa gtggtctcag agttgagtta ccttggcttc tgggagggtga 1260
aactgtatcc ctataccctg aagctttaag ggggtgcaat gtagatgaga cccaacata 1320
gacctcttc acag 1334

<210> 45

<211> 512

<212> DNA

<213> HUMAN

<220>

<221> intron

<222> (1)..(512)

<223> 7th MN intron

<400> 45

gtgggcctgg ggtgtgtgtg gacacagtgg gtgcggggga aagaggatgt aagatgagat 60

gagaaacagg agaagaaaga aatcaaggct gggctctgtg gcttacgcct ataatcccac 120

cacgttggga ggctgaggtg ggagaatggt ttgagcccag gagttcaaga caaggcgggg 180

caacatagtg tgaccccatc tctaccaaaa aaaccccaac aaaacaaaaa atagccgggc 240

atggtggtat ggggcctagt cccagctact caaggaggct gaggtgggaa gatcgcttga 300

ttccaggagt ttgagactgc agtgagctat gatcccacca ctgcctacca tctttaggat 360

acattttattt atttataaaa gaaatcaaga ggctggatgg ggaatacagg agctggaggg 420

tggagccctg aggtgtgtgt tgtgagctgg cctggggacc ttgtttcctg tcatgccatg 480

aaccacacca cactgtccac tgacctcct ag 512

<210> 46

<211> 114

<212> DNA

<213> HUMAN

<220>

<221> intron

<222> (1)..(114)

<223> 8th MN intron

<400> 46

gtacagcttt gtctggtttc cccccagcca gtagtcctt atcctcccat gtgtgtgcc 60

gtgtctgtca ttggtggtca cagcccgct ctcacatctc ctttttctct ccag 114

<210> 47

<211> 617

<212> DNA

<213> HUMAN

<220>

<221> intron

<222> (1)..(617)

<223> 9th MN intron

<400> 47

gtgagtctgc ccctcctctt ggtcctgatg ccaggagact cctcagcacc attcagcccc 60
agggctgctc aggaccgcct ctgctccctc tcctttttctg cagaacagac cccaacccca 120
atattagaga ggcagatcat ggtgggggatt cccccattgt cccagagggc taattgatta 180
gaatgaagct tgagaaatct cccagcatcc ctctcgcaaa agaatcccc cccctttttt 240
taaagatagg gtctcactct gtttgcccca ggctgggggtg ttgtggcacg atcatagctc 300
actgcagcct cgaactccta ggctcaggca atcctttcac cttagcttct caaagcactg 360
ggactgtagg catgagccac tgtgcctggc cccaaacggc ccttttactt ggcttttagg 420
aagcaaaaac ggtgcttata ttacccttc tcgtgtatcc accctcatcc cttggctggc 480
ctctttctgga gactgaggca ctatggggct gcctgagaac tcggggcagg ggtgggtggag 540
tgcactgagg cagggtgtga ggaactctgc agaccctct tccttcccaa agcagccctc 600
tctgctctcc atcgag 617

<210> 48

<211> 130

<212> DNA

<213> HUMAN

<220>

<221> intron

<222> (1)..(130)

<223> 10th MN intron

<400> 48

gtattacact gaccctttct tcaggcacia gttccccca cccttgtgga gtcacttcat 60
gcaaagcgca tgcaaatgag ctgctcctgg gccagttttc tgattagcct ttctgtttgt 120
gtacacacag 130

<210> 49

<211> 1401

<212> DNA

<213> HUMAN

<400> 49

caaaactttca cttttgttgc ccaggctgga gtgcaatggc gcgatctcgg ctcaactgcaa 60
cctccacctc ccgggttcaa gtgattctcc tgcctcagcc tctagccaag tagctgcgat 120
tacaggcatg cgccaccacg cccggctaatt ttttgtattt ttagtagaga cgggggtttcg 180
ccatgttggt caggctggtc tcgaactcct gatctcaggt gatccaacca ccctggcctc 240
ccaaagtgct gggattatag gcgtgagcca cagcgcctgg cctgaagcag ccactcactt 300
ttacagaccc taagacaatg attgcaagct ggtaggattg ctgtttggcc caccacagctg 360
cgggtgttgag tttgggtgcg gtctcctgtg ctttgcacct ggcccgtta aggcatttgt 420
tacccgtaat gtcctgtaa ggcactctcg tttgtgacat cgttttggtc gccaggaagg 480
gattggggct ctaagcttga gcggttcac cttttcattt atacagggga tgaccagagt 540
cattggcgt atggaggta gacaccacc cgctgcacag acccaatctg ggaaccacg 600
tctgtggatc tcccctacag ccgtccctga aactgggtcc cgggcgtccc acccgccgcc 660
caccgtcca cccctcacc ttttctacc gggttcccta agttcctgac ctaggcgtca 720
gacttctca ctatactctc ccaccccagg cgaccgcc tggccccggg tgtccccagc 780
ctgcgcgggc cgcttcagtc ccccggtgga tatccgcccc cagctcgccg cttctgccc 840
ggccctgcgc cccctggaac tcctgggctt ccagctcccg ccgctcccag aactgcgcct 900
gcgcaacaat ggccacagt gtgaggggggt ctccccgccg agacttgggg atggggcggg 960
gcgcagggaa gggaaccgtc gcgcagtgc tgcctggggg ttgggctggc cctaccgggc 1020
ggggccggct cacttgctc tccctacgca gtgcaactga ccctgcctcc tgggctagag 1080
atggctctgg gtccccggcg ggagtaccgg gctctgcagc tgcacttgca ctggggggct 1140
gcaggtcgtc cgggctcgga gcacactgtg gaaggccacc gttccctgc cgaggtgagc 1200
gcggactggc cgagaagggg caaaggagcg gggcggacgg gggccagaga cgtggccctc 1260
tcctaccctc gtgtcctttt cagatccacg tggttcacct cagcaccgcc tttgccagag 1320

ttgacgagggc cttgggggcgc ccgggaggcc tggccgtggtt ggccgccttt ctggagggtac 1380

cagatcctgg acaccccccta c 1401

<210> 50

<211> 59

<212> PRT

<213> HUMAN

<400> 50

Ser Ser Gly Glu Asp Asp Pro Leu Gly Glu Glu Asp Leu Pro Ser Glu
1 5 10 15

Glu Asp Ser Pro Arg Glu Glu Asp Pro Pro Gly Glu Glu Asp Leu Pro
20 25 30

Gly Glu Glu Asp Leu Pro Gly Glu Glu Asp Leu Pro Glu Val Lys Pro
35 40 45

Lys Ser Glu Glu Glu Gly Ser Leu Lys Leu Glu
50 55

<210> 51

<211> 257

<212> PRT

<213> HUMAN

<400> 51

Gly Asp Asp Gln Ser His Trp Arg Tyr Gly Gly Asp Pro Pro Trp Pro
1 5 10 15

Arg Val Ser Pro Ala Cys Ala Gly Arg Phe Gln Ser Pro Val Asp Ile
20 25 30

Arg Pro Gln Leu Ala Ala Phe Cys Pro Ala Leu Arg Pro Leu Glu Leu
35 40 45

Leu Gly Phe Gln Leu Pro Pro Leu Pro Glu Leu Arg Leu Arg Asn Asn
50 55 60

Gly His Ser Val Gln Leu Thr Leu Pro Pro Gly Leu Glu Met Ala Leu
65 70 75 80

Gly Pro Gly Arg Glu Tyr Arg Ala Leu Gln Leu His Leu His Trp Gly
85 90 95

Ala Ala Gly Arg Pro Gly Ser Glu His Thr Val Glu Gly His Arg Phe
 100 105 110

Pro Ala Glu Ile His Val Val His Leu Ser Thr Ala Phe Ala Arg Val
 115 120 125

Asp Glu Ala Leu Gly Arg Pro Gly Gly Leu Ala Val Leu Ala Ala Phe
 130 135 140

Leu Glu Glu Gly Pro Glu Glu Asn Ser Ala Tyr Glu Gln Leu Leu Ser
 145 150 155 160

Arg Leu Glu Glu Ile Ala Glu Glu Gly Ser Glu Thr Gln Val Pro Gly
 165 170 175

Leu Asp Ile Ser Ala Leu Leu Pro Ser Asp Phe Ser Arg Tyr Phe Gln
 180 185 190

Tyr Glu Gly Ser Leu Thr Thr Pro Pro Cys Ala Gln Gly Val Ile Trp
 195 200 205

Thr Val Phe Asn Gln Thr Val Met Leu Ser Ala Lys Gln Leu His Thr
 210 215 220

Leu Ser Asp Thr Leu Trp Gly Pro Gly Asp Ser Arg Leu Gln Leu Asn
 225 230 235 240

Phe Arg Ala Thr Gln Pro Leu Asn Gly Arg Val Ile Glu Ala Ser Phe
 245 250 255

Pro

<210> 52

<211> 20

<212> PRT

<213> HUMAN

<400> 52

Ile Leu Ala Leu Val Phe Gly Leu Leu Phe Ala Val Thr Ser Val Ala
 1 5 10 15

Phe Leu Val Gln
 20

<210> 53
 <211> 25
 <212> PRT
 <213> HUMAN

<400> 53

Met Arg Arg Gln His Arg Arg Gly Thr Lys Gly Gly Val Ser Tyr Arg
 1 5 10 15

Pro Ala Glu Val Ala Glu Thr Gly Ala
 20 25

<210> 54
 <211> 59
 <212> PRT
 <213> HUMAN

<400> 54

Ser Ala Ser Glu Glu Pro Ser Pro Ser Glu Val Pro Phe Pro Ser Glu
 1 5 10 15

Glu Pro Ser Pro Ser Glu Glu Pro Phe Pro Ser Val Arg Pro Phe Pro
 20 25 30

Ser Val Val Leu Phe Pro Ser Glu Glu Pro Phe Pro Ser Lys Glu Pro
 35 40 45

Ser Pro Ser Glu Glu Pro Ser Ala Ser Glu Glu
 50 55

<210> 55
 <211> 470
 <212> RNA
 <213> HUMAN

<400> 55

cauggcccccg auaaccuucu gccugugcac acaccugccc cucacuccac ccccauccua 60
 gcuuugguau gggggagagg gcacagggcc agacaaaccu gugagacuuu ggcuccaucu 120
 cugcaaaagg ggcgucugug agucagccug cuccccucca ggcuugcucc uccccaccc 180
 agcucucguu uccaaugcac guacagcccc uacacaccgu gugcugggac accccacagu 240
 cagccgcaug gcuccccugu gccccagccc cuggcucccu cuguugauc cggccccugc 300

uccaggccuc acugugcaac ugcugcuguc acugcugcuu cuggugccug uccaucceca 360
gagguugccc cggaugcagg aggauuccecu cuugggagga ggcucuucug gggaagauga 420
cccacugggc gaggaggauc ugcccaguga agaggauuca cccagagagg 470

<210> 56
<211> 292
<212> DNA
<213> HUMAN

<400> 56
gtttttttga gacggagtct tgcattctgtc atgcccaggc tggagtagca gtggtgcat 60
ctcggctcac tgcaagctcc acctcccgag ttcacgcat tttcctgcct cagcctcccg 120
agtagctggg actacaggcg cccgccacca tgcccggcta attttttgta tttttgtag 180
agacgggggt tcaccgtgtt agccagaatg gtctcgatct cctgacttcg tgatccacc 240
gcctcggcct cccaaagtcc tgggattaca ggtgtgagcc accgcacctg gc 292

<210> 57
<211> 262
<212> DNA
<213> HUMAN

<400> 57
tttttttttt gagacagggt cttgctctgt caccaggcc agagtgcaat ggtacagtct 60
cagctcactg cagcctcaac cgcctcggt caaacatca tcccatttca gcctcctgag 120
tagctgggac tacaggcaca tgccattaca cctggctaat tttttgtat ttctagtaga 180
gacagggttt ggccatgttg cccgggctgg tctcgaactc ctggactcaa gcaatccacc 240
cacctcagcc tcccataatg ag 262

<210> 58
<211> 2501
<212> DNA
<213> HUMAN

<220>
<221> misc_feature

<222> (1) .. (2501)

<400> 58

tggtgactcg tgaccttacc cccaaccctg tgctctctga aacatgagct gtgtccactc 60
agggttaaataa ggattaaggg cggtgcaaga tgtgctttgt taaacagatg cttgaaggca 120
gcatgctcgt taagagtcac caccaatccc taatctcaag taatcaggga cacaaacact 180
gcggaaggcc gcagggtcct ctgcctagga aaaccagaga cctttgttca cttgtttatc 240
tgaccttccc tccactattg tccatgacct tgccaaatcc cctctgtga gaaacaccca 300
agaattatca ataaaaaaat aaatttaaaa aaaaaatata aaaaaaaaaa aaaaaaaaaa 360
aaaagactta cgaatagtta ttgataaatg aatagctatt ggtaaagcca agtaaagtat 420
catattcaaa accagacggc catcatcaca gctcaagtct acctgatttg atctctttat 480
cattgtcatt ctttggtatc actagattag tcatcatcct caaaattctc cccaagttc 540
taattacgtt ccaaacattt aggggttaca tgaagcttga acctactacc ttctttgctt 600
ttgagccatg agttgtagga atgatgagtt tacaccttac atgctgggga ttaatttaaa 660
ctttacctct aagtcagttg ggtagccttt ggcttatttt tgtagctaata tttgtagtta 720
atggatgcac tgtgaatctt gctatgatag ttttcctcca cactttgcca ctaggggtag 780
gtaggtactc agttttcagt aattgcttac ctaagacctt aagccctatt tctctgttac 840
tggcctttat ctgtaatatg ggcatattta atacaatata atttttggag tttttttggt 900
tgtttggttg tttgtttttt tgagacggag tcttgcatct gtcatgcca ggctggagta 960
gcagtgggtc catctcggct cactgcaagc tccacctccc gagttcacgc cattttcctg 1020
cctcagcctc ccgagtagct gggactacag gcgcccgcga ccatgcccgg ctaatttttt 1080
gtatttttgg tagagacggg gtttcacctg gtttagccaga atggtctcga tctcctgact 1140
tcgtgatcca ccgcctcgg cctcccaaag ttctgggatt acaggtgtga gccaccgcac 1200
ctggccaatt ttttgagtct tttaaagtaa aaatatgtct tgtaagctgg taactatggt 1260
acatttcctt ttattaatgt ggtgctgacg gtcatatagg ttcttttgag tttggcatgc 1320
atatgctact ttttgacgtc ctttcattac atttttctct cttcatttga agagcatggt 1380

atatctttta gcttcacttg gcttaaaagg ttctctcatt agcctaacac agtgtcattg 1440
ttggtaccac ttggatcata agtggaaaaa cagtcaagaa attgcacagt aatacttgtt 1500
tgtaagaggg atgattcagg tgaatctgac actaagaaac tcccctacct gaggtctgag 1560
attcctctga cattgctgta tataggcttt tcctttgaca gcctgtgact gcggactatt 1620
tttcttaagc aagatatgct aaagttttgt gagccttttt ccagagagag gtctcatatc 1680
tgcacatcaagt gagaacatat aatgtctgca tgtttccata tttcaggaat gtttgcttgt 1740
gttttatgct tttatataga cagggaaact tgttcctcag tgacccaaaa gaggtgggaa 1800
ttgttattgg atatcatcat tggcccacgc tttctgacct tggaaacaat taagggttca 1860
taatctcaat tctgtcagaa ttggtacaag aaatagctgc tatgtttctt gacattccac 1920
ttggtaggaa ataagaatgt gaaactcttc agttgggtgtg tgtccctngt ttttttgcaa 1980
tttctttctt actgtgttaa aaaaaagtat gatcttgctc tgagaggtga ggcattctta 2040
atcatgatct ttaaagatca ataataaat cctttcaagg attatgtctt tattataata 2100
aagataatth gtctttaaca gaatcaataa tataatccct taaaggatta tatctttgct 2160
gggcgcagtg gctcacacct gtaatcccag cactttgggt ggccaagggtg gaaggatcaa 2220
atttgcttac ttctatatta tcttctaaag cagaattcat ctctcttccc tcaatatgat 2280
gatattgaca gggtttgccc tcaactacta gattgtgagc tcctgctcag ggcaggtagc 2340
gttttttgtt tttgtttttg tttttctttt ttgagacagg gtcttgctct gtcaccacagg 2400
ccagagtgca atggtacagt ctcagctcac tgcagcctca accgcctcgg ctcaaaccat 2460
catcccattt cagcctcctg agtagctggg actacaggca c 2501

<210> 59

<211> 292

<212> DNA

<213> HUMAN

<220>

<221> misc_feature

<222> (1)

<400> 59

ttttttttgag acggagtctt gcatctgtca tgcccaggct ggagtagcag tgggtgccatc 60
 teggctcact gcaagctcca cctcccgagt tcacgccatt ttctgcctc agcctcccg 120
 gtagctggga ctacaggcgc ccgccaccat gcccggttaa ttttttgtat ttttggtaga 180
 gacgggggttt caccgtgtta gccagaatgg tctcgatctc ctgacttcgt gatccacccg 240
 cctcggcctc ccaaagttct gggattacag gtgtgagcca ccgcacctgg cc 292

<210> 60

<211> 262

<212> DNA

<213> HUMAN

<400> 60

ttcttttttg agacagggtc ttgctctgtc acccaggcca gaggcaatg gtacagtctc 60
 agctcactgc agcctcaacc gcctcggctc aaaccatcat cccatttcag cctcctgagt 120
 agctgggact acaggcacat gccattacac ctggctaatt tttttgtatt tctagtagag 180
 acagggtttg gccatgttgc ccgggctggt ctogaactcc tggactcaag caatccaccc 240
 acctcagcct cccaaaatga gg 262

<210> 61

<211> 294

<212> DNA

<213> HUMAN

<400> 61

tttttttttg agacaaactt tcaacttttg tgcccaggct ggagtgcaat ggcgcgatct 60
 cggctcactg caacctccac ctcccgggtt caagtgatc ttctgcctca gcctctagcc 120
 aagtagctgc gattacaggc atgcgccacc acgcccggct aatttttgta ttttagtag 180
 agacgggggtt tcgcatgtt ggtcaggctg gtctcgaact cctgatctca ggtgatccaa 240
 ccaccctggc ctcccaaagt gctgggatta taggcgtgag ccacagcgcc tggc 294

<210> 62

<211> 276

<212> DNA

<213> HUMAN

<400> 62

tgacagtctc tctgtcgccc aggctggagt gcagtgggtg gatcttgggt cactgcaact 60
tccgcctccc gggttcaagg gattctcctg cctcagcttc ctgagtagct ggggttacag 120
gtgtgtgcc aatgcccag ctaatttttt tttgtatttt tagtagacag ggtttcacca 180
tgttggtcag gctgggtctca aactcctggc ctcaagtgat ccgcctgact cagcctacca 240
aagtgtgat tacaagtgtg agccaccgtg ccagc 276

<210> 63

<211> 289

<212> DNA

<213> HUMAN

<400> 63

cgccgggcac ggtgggtcac gcctgtaatc ccagcacttt gggaggccaa ggcagggtgga 60
tcacgaggtc aagagatcaa gaccatcctg gccaacatgg tgaaacccca tctctactaa 120
aaatacgaaa aaatagccag gcgtgggtggc ggggtgcctgt aatcccagct actcgggagg 180
ctgaggcagg agaatggcat gaaccggga ggcagaagtt gcagtgagcc gagatcgtgc 240
cactgcactc cagcctgggc aacagagcga gactcttgct tcaaaaaaa 289

<210> 64

<211> 298

<212> DNA

<213> HUMAN

<400> 64

aggctgggct ctgtggctta cgcctataat cccaccacgt tgggaggctg aggtgggaga 60
atggtttgag ccagaggtt caagacaagg cggggcaaca tagtgtgacc ccattctctac 120
caaaaaaacc ccaacaaaac caaaaatagc cgggcatggt ggtatgcggc ctatgccag 180
ctactcaagg aggctgaggt gggaagatcg cttgattcca ggagtttgag actgcagtga 240
gctatgatcc caccactgcc taccatcttt aggatacatt tattttattta taaaagaa 298

<210> 65
<211> 105
<212> DNA
<213> HUMAN

<400> 65
ttttttacat ctttagtaga gacagggttt caccatattg gccaggctgc tctcaaactc 60
ctgaccttgt gatccaccag cctcggcctc ccaaagtgt gggat 105

<210> 66
<211> 83
<212> DNA
<213> HUMAN

<400> 66
cctcgaactc ctaggctcag gcaatccttt caccttagct tctcaaagca ctgggactgt 60
aggcatgagc cactgtgcct ggc 83

<210> 67
<211> 11
<212> DNA
<213> HUMAN

<400> 67
agaaggtaag t 11

<210> 68
<211> 11
<212> DNA
<213> HUMAN

<400> 68
tggagggtgag a 11

<210> 69
<211> 11
<212> DNA
<213> HUMAN

<400> 69

cagtcgtgag g

11

<210> 70

<211> 11

<212> DNA

<213> HUMAN

<400> 70

ccgaggtgag c

11

<210> 71

<211> 11

<212> DNA

<213> HUMAN

<400> 71

tggaggtacc a

11

<210> 72

<211> 11

<212> DNA

<213> HUMAN

<400> 72

ggaaggtcag t

11

<210> 73

<211> 11

<212> DNA

<213> HUMAN

<400> 73

agcaggtggg c

11

<210> 74

<211> 11

<212> DNA

<213> HUMAN

<400> 74

gccaggtaca g

11

<210> 75

<211> 11

<212> DNA

<213> HUMAN

<400> 75

tgctggtgag t

11

<210> 76

<211> 11

<212> DNA

<213> HUMAN

<400> 76

atacagggga t

11

<210> 77

<211> 11

<212> DNA

<213> HUMAN

<400> 77

atacagggga t

11

<210> 78

<211> 11

<212> DNA

<213> HUMAN

<400> 78

ccccaggcga c

11

<210> 79

<211> 11

<212> DNA

<213> HUMAN

<400> 79

acgcagtgca a

11

<210> 80

<211> 11

<212> DNA

<213> HUMAN

<400> 80

tttcagatcc a

11

<210> 81

<211> 11

<212> DNA

<213> HUMAN

<400> 81

ccccaggagg g

11

<210> 82

<211> 11

<212> DNA

<213> HUMAN

<400> 82

tcacaggctc a

11

<210> 83

<211> 11

<212> DNA

<213> HUMAN

<400> 83

ccctagctcc a

11

<210> 84

<211> 11

<212> DNA

<213> HUMAN

<400> 84

ctccagtcca g

11

<210> 85

<211> 12

<212> DNA

<213> HUMAN

<400> 85

tcgcaggtga ca

12

<210> 86

<211> 11

<212> DNA

<213> HUMAN

<400> 86

acacagaagg g

11

<210> 87

<211> 377

<212> PRT

<213> HUMAN

<400> 87

Gln Arg Leu Pro Arg Met Gln Glu Asp Ser Pro Leu Gly Gly Gly Ser
 1 5 10 15

Ser Gly Glu Asp Asp Pro Leu Gly Glu Glu Asp Leu Pro Ser Glu Glu
 20 25 30

Asp Ser Pro Arg Glu Glu Asp Pro Pro Gly Glu Glu Asp Leu Pro Gly
 35 40 45

Glu Glu Asp Leu Pro Gly Glu Glu Asp Leu Pro Glu Val Lys Pro Lys
 50 55 60

Ser Glu Glu Glu Gly Ser Leu Lys Leu Glu Asp Leu Pro Thr Val Glu
 65 70 75 80

Ala Pro Gly Asp Pro Gln Glu Pro Gln Asn Asn Ala His Arg Asp Lys
 85 90 95

Glu Gly Asp Asp Gln Ser His Trp Arg Tyr Gly Gly Asp Pro Pro Trp
 100 105 110

Pro Arg Val Ser Pro Ala Cys Ala Gly Arg Phe Gln Ser Pro Val Asp
 115 120 125

Ile Arg Pro Gln Leu Ala Ala Phe Cys Pro Ala Leu Arg Pro Leu Glu
 130 135 140

Leu Leu Gly Phe Gln Leu Pro Pro Leu Pro Glu Leu Arg Leu Arg Asn
 145 150 155 160

Asn Gly His Ser Val Gln Leu Thr Leu Pro Pro Gly Leu Glu Met Ala
 165 170 175
 Leu Gly Pro Gly Arg Glu Tyr Arg Ala Leu Gln Leu His Leu His Trp
 180 185 190
 Gly Ala Ala Gly Arg Pro Gly Ser Glu His Thr Val Glu Gly His Arg
 195 200 205
 Phe Pro Ala Glu Ile His Val Val His Leu Ser Thr Ala Phe Ala Arg
 210 215 220
 Val Asp Glu Ala Leu Gly Arg Pro Gly Gly Leu Ala Val Leu Ala Ala
 225 230 235 240
 Phe Leu Glu Glu Gly Pro Glu Glu Asn Ser Ala Tyr Glu Gln Leu Leu
 245 250 255
 Ser Arg Leu Glu Glu Ile Ala Glu Glu Gly Ser Glu Thr Gln Val Pro
 260 265 270
 Gly Leu Asp Ile Ser Ala Leu Leu Pro Ser Asp Phe Ser Arg Tyr Phe
 275 280 285
 Gln Tyr Glu Gly Ser Leu Thr Thr Pro Pro Cys Ala Gln Gly Val Ile
 290 295 300
 Trp Thr Val Phe Asn Gln Thr Val Met Leu Ser Ala Lys Gln Leu His
 305 310 315 320
 Thr Leu Ser Asp Thr Leu Trp Gly Pro Gly Asp Ser Arg Leu Gln Leu
 325 330 335
 Asn Phe Arg Ala Thr Gln Pro Leu Asn Gly Arg Val Ile Glu Ala Ser
 340 345 350
 Phe Pro Ala Gly Val Asp Ser Ser Pro Arg Ala Ala Glu Pro Val Gln
 355 360 365
 Leu Asn Ser Cys Leu Ala Ala Gly Asp
 370 375

<210> 88

<211> 34

<212> DNA

<213> HUMAN

<400> 88

tagacagatc tacgatggct cccctgtgcc ccag

34

<210> 89

<211> 34

<212> DNA

<213> HUMAN

<400> 89

attcctctag acagttaccg gctccccctc agat

34

<210> 90

<211> 3532

<212> DNA

<213> HUMAN

<220>

<221> misc_feature

<222> (1)..(3532)

<400> 90

tgttgactcg tgaccttacc cccaaccctg tgctctctga aacatgagct gtgtccactc 60

agggttaaat ggattaaggg cgggtgcaaga tgtgctttgt taaacagatg cttgaaggca 120

gcatgctcgt taagagtcac caccaatccc taatctcaag taatcaggga cacaaacact 180

gcggaaggcc gcagggtcct ctgcctagga aaaccagaga cctttgttca cttgtttatc 240

tgaccttccc tccactattg tccatgaccc tgccaaatcc ccctctgtga gaaacaccca 300

agaattatca ataaaaaaat aaatttaaaa aaaaaataca aaaaaaaaaa aaaaaaaaaa 360

aaaagactta cgaatagtta ttgataaatg aatagctatt ggtaaagcca agtaaagatg 420

catattcaaa accagacggc catcatcaca gctcaagtct acctgatttg atctctttat 480

cattgtcatt ctttggtatc actagattag tcatcatcct caaaattctc cccaagtgc 540

taattacgtt ccaaacattt aggggttaca tgaagcttga acctactacc ttctttgctt 600

ttgagccatg agttgtagga atgatgagtt tacaccttac atgctgggga ttaatttaaa 660

ctttacctct aagtcagttg ggtagccttt ggcttatttt tgtagctaatt tttgtagtta 720

atggatgcac tgtgaatcct gctatgatag ttttctcca cactttgcca ctaggggtag 780

gtaggtactc agttttcagt aattgcttac ctaagaccct aagccctatt tctcttgtac 840
tggcctttat ctgtaatatg ggcataatata atacaatata attttttgag tttttttgtt 900
tgtttggttg tttgtttttt tgagacggag tcttgcatct gtcatgccca ggctggagta 960
gcagtgggtgc catctcggct cactgcaagc tccacctccc gagttcacgc cattttcctg 1020
cctcagcctc ccgagtagct gggactacag gcgcccgcga ccatgcccgg ctaatttttt 1080
gtattttttg tagagacggg gtttcaccgt gtttagccaga atggctctga tctcctgact 1140
tcgtgatcca cccgcctcgg cctcccaaag ttctgggatt acaggtgtga gccaccgcac 1200
ctggccaatt ttttgagtct tttaaagtaa aaatatgtct tgtaagctgg taactatggt 1260
acatttcctt ttattaatgt ggtgctgacg gtcatatagg ttcttttgag tttggcatgc 1320
atatgctact ttttgcatc ctttcattac atttttctct cttcatttga agagcatggt 1380
atatctttta gcttcacttg gcttaaaagg ttctctcatt agcctaacac agtgtcattg 1440
ttggtaccac ttggatcata agtggaaaaa cagtcaagaa attgcacagt aatacttgtt 1500
tgtaagaggg atgattcagg tgaatctgac actaagaaac tcccctacct gaggtctgag 1560
attcctctga cattgctgta tataggcttt tcctttgaca gcctgtgact gcggactatt 1620
tttcttaagc aagatatgct aaagttttgt gagccttttt ccagagagag gtctcatatc 1680
tgcatcaagt gagaacatat aatgtctgca tgtttcata tttcaggaat gtttgcttgt 1740
gttttatgct tttatataga cagggaaact tgttcctcag tgacccaaaa gaggtgggaa 1800
ttgttattgg atatcatcat tggcccacgc tttctgacct tggaaacaat taagggttca 1860
taatctcaat tctgtcagaa ttggtacaag aaatagctgc tatgtttctt gacattccac 1920
ttggtaggaa ataagaatgt gaaactcttc agttgggtgtg tgtccctngt ttttttgcaa 1980
tttccttctt actgtgttaa aaaaaagtat gatcttgctc tgagaggtga ggcattctta 2040
atcatgatct ttaaagatca ataataat ctttcaagg attatgtctt tattataata 2100
aagataattt gtctttaaca gaatcaataa tataatccct taaaggatta tatctttgct 2160
gggcgcagtg gctcacacct gtaatcccag cactttgggt ggccaagggt gaaggatcaa 2220

atttgectac ttctatatta tcttctaaag cagaattcat ctctcttccc tcaatatgat 2280
gatattgaca gggtttgccc tcactcacta gattgtgagc tcctgctcag ggcaggtagc 2340
gtttttttgtt tttgtttttg tttttctttt ttgagacagg gtcttgctct gtcacccagg 2400
ccagagtgca atggtacagt ctgagctcac tgcagcctca accgcctcgg ctcaaaccat 2460
catcccattt cagcctcctg agtagctggg actacaggca catgccatta cacctggcta 2520
atttttttgt atttctagta gagacagggt ttggccatgt tgcccgggct ggtctcgaac 2580
tcctggactc aagcaatcca cccacctcag cctcccaaaa tgagggaccg tgtcttattc 2640
atttccatgt ccctagtcca tagcccagtg ctggacctat ggtagtacta aataaatatt 2700
tgttgaatgc aatagtaa atgcatttcag ggagcaagaa ctagattaac aaaggtggta 2760
aaaggtttgg agaaaaaat aatagtttaa tttggctaga gtatgaggga gagtagtagg 2820
agacaagatg gaaaggtctc ttgggcaagg ttttgaagga agttggaagt cagaagtaca 2880
caatgtgcat atcgtggcag gcagtgggga gccaatgaag gctttttgagc aggagagtaa 2940
tgtgttgaaa aataaatata ggttaaacct atcagagccc ctctgacaca tacacttgct 3000
tttcattcaa gctcaagttt gtctcccaca taccattac ttaactcacc ctcgggctcc 3060
cctagcagcc tgccctacct ctttacctgc ttcttggtgg agtcagggat gtatacatga 3120
gctgctttcc ctctcagcca gaggacatgg ggggccccag ctcccctgcc tttccccttc 3180
tgtgcctgga gctgggaagc aggccagggt tagctgaggc tggctggcaa gcagctgggt 3240
ggtgccaggg agagcctgca tagtgccagg tggcgcttg ggttccaagc tagtccatgg 3300
ccccgataac cttctgcctg tgcacacacc tgcccctcac tccaccccca tcctagcttt 3360
ggtatggggg agagggcaca gggccagaca aacctgtgag actttggctc catctctgca 3420
aaagggcgct ctgtgagtca gcctgctccc ctccaggctt gctcctcccc caccagctc 3480
tcgtttccaa tgcacgtaca gcccgtagac accgtgtgct gggacacccc ac 3532

<210> 91

<211> 204

<212> DNA

<213> HUMAN

<400> 91

cctgccccctc actccacccc catcctagct ttggtatggg ggagagggca cagggccaga 60

caaacctgtg agactttggc tccatctctg caaaagggcg ctctgtgagt cagcctgctc 120

ccctccaggc ttgctcctcc cccaccagc tctcgtttcc aatgcacgta cagcccgtac 180

acaccgtgtg ctgggacacc ccac 204

<210> 92

<211> 132

<212> DNA

<213> HUMAN

<400> 92

ggatcctgtt gactcgtgac cttaccccca accctgtgct ctctgaaaca tgagctgtgt 60

ccactcaggg ttaaattgat taagggcggg gcaagatgtg ctttggttaa cagatgcttg 120

aaggcagcat gc 132

<210> 93

<211> 275

<212> DNA

<213> HUMAN

<400> 93

gcatagtgcc aggtggtgcc ttgggttcca agctagtcca tggccccgat aaccttctgc 60

ctgtgcacac acctgcccct cactccaccc ccatoctagc ttggtatgg gggagagggc 120

acagggccag acaaacctgt gagactttgg ctccatctct gcaaaagggc gctctgtgag 180

tcagcctgct cccctccagg cttgctcctc ccccaccag ctctcgtttc caatgcacgt 240

acagcccgtc cacaccgtgt gctgggacac cccac 275

<210> 94

<211> 89

<212> DNA

<213> HUMAN

<400> 94

ctgctccccct ccaggcttgc tcctccccc cccagctctc gtttccaatg cacgtacagc 60

ccgtacacac cgtgtgctgg gacacccca

89

<210> 95

<211> 61

<212> DNA

<213> HUMAN

<400> 95

caccagctc tcgtttccaa tgcacgtaca gcccgtagac accgtgtgct gggacacccc 60

a

61

<210> 96

<211> 116

<212> DNA

<213> HUMAN

<400> 96

acctgccccct cactccaccc ccatactagc tttggtatgg gggagagggc acagggccag 60

acaaacctgt gagactttgg ctccatctct gcaaaagggc gctctgtgag tcagcc 116

<210> 97

<211> 36

<212> PRT

<213> HUMAN

<400> 97

Gly Glu Glu Asp Leu Pro Ser Glu Glu Asp Ser Pro Arg Glu Glu Asp

1

5

10

15

Pro Pro Gly Glu Glu Asp Leu Pro Gly Glu Glu Asp Leu Pro Gly Glu

20

25

30

Glu Asp Leu Pro

35

<210> 98

<211> 6

<212> PRT

<213> HUMAN

<400> 98

Gly Glu Glu Asp Leu Pro

1

5

<210> 99

<211> 4

<212> PRT

<213> HUMAN

<400> 99

Glu Glu Asp Leu

1

<210> 100

<211> 5

<212> PRT

<213> HUMAN

<400> 100

Glu Glu Asp Leu Pro

1

5

<210> 101

<211> 6

<212> PRT

<213> HUMAN

<400> 101

Glu Asp Leu Pro Ser Glu

1

5

<210> 102

<211> 7

<212> PRT

<213> HUMAN

<400> 102

Glu Glu Asp Leu Pro Ser Glu

1

5

<210> 103

<211> 6

<212> PRT

<213> HUMAN

<400> 103

Asp Leu Pro Gly Glu Glu

1

5

<210> 104

<211> 22

<212> PRT

<213> HUMAN

<400> 104

Gly Gly Ser Ser Gly Glu Asp Asp Pro Leu Gly Glu Glu Asp Leu Pro

1

5

10

15

Ser Glu Glu Asp Ser Pro

20

<210> 105

<211> 25

<212> PRT

<213> HUMAN

<400> 105

Gly Glu Glu Asp Leu Pro Ser Glu Glu Asp Ser Pro Arg Glu Glu Asp

1

5

10

15

Pro Pro Gly Glu Glu Asp Leu Pro Gly

20

25

<210> 106

<211> 24

<212> PRT

<213> HUMAN

<400> 106

Glu Asp Pro Pro Gly Glu Glu Asp Leu Pro Gly Glu Glu Asp Leu Pro

1

5

10

15

Gly Glu Glu Asp Leu Pro Glu Val

20

<210> 107

<211> 7

<212> PRT

<213> HUMAN

<400> 107

Gly Glu Thr Arg Ala Pro Leu

1

5

<210> 108

<211> 7

<212> PRT

<213> HUMAN

<400> 108

Gly Glu Thr Arg Glu Pro Leu

1

5

<210> 109

<211> 7

<212> PRT

<213> HUMAN

<400> 109

Gly Gln Thr Arg Ser Pro Leu

1

5

<210> 110

<211> 1247

<212> DNA

<213> HUMAN

<220>

<221> misc_feature

<222> (1)..(1247)

<400> 110

tatgctactt tttgcagtc tttcattaca tttttctctc ttcatttgaa gagcatgtta 60

tatcttttag cttcacttgg cttaaaagggt totctcatta gcctaacaca gtgtcattgt 120

tggtaccact tggatcataa gtggaaaaaac agtcaagaaa ttgcacagta atacttgttt 180

gtaagaggga tgattcaggt gaatctgaca ctaagaaact cccctacctg aggtctgaga 240

ttcctctgac attgctgtat ataggctttt cctttgacag cctgtgactg cggactattt 300

ttcttaagca agatatgcta aagttttgtg agcctttttc cagagagagg tctcatatct 360
 gcatcaagtg agaacatata atgtctgcat gtttccatat ttcaggaatg tttgcttgtg 420
 ttttatgctt ttatatagac agggaaactt gttcctcagt gacccaaaag aggtgggaat 480
 tgttattgga tatcatcatt ggcccacgct ttctgacctt ggaaacaatt aagggttcat 540
 aatctcaatt ctgtcagaat tgggtacaaga aatagctgct atgtttcttg acattccact 600
 tggtaggaaa taagaatgtg aaactcttca gttgggtgtg gtcctngtt tttttgcaat 660
 ttccttctta ctgtgttaaa aaaaagtatg atcttgctct gagaggtgag gcattcttaa 720
 tcatgatctt taaagatcaa taatataatc ctttcaagga ttatgtcttt attataataa 780
 agataatttg tctttaacag aatcaataat ataatccctt aaaggattat atctttgctg 840
 ggcgagtggtg ctcacacctg taatcccagc actttgggtg gccaaagggtg aaggatcaaa 900
 tttgcctact tctatattat cttctaaagc agaattcatc tctcttcctt caatatgatg 960
 atattgacag ggtttgcctt cactcactag attgtgagct cctgctcagg gcaggtagcg 1020
 ttttttgttt ttgtttttgt ttttcttttt tgagacaggg tcttgctctg tcaccagggc 1080
 cagagtgcaa tgggtacagtc tcagctcact gcagcctcaa ccgcctcggc tcaaaccatc 1140
 atcccatttc agcctcctga gtagctggga ctacaggcac atgccattac acctgggctaa 1200
 tttttttgta tttctagtag agacaggggtt tggccatggt gcccggg 1247

<210> 111

<211> 17

<212> DNA

<213> HUMAN

<400> 111

ctctgtgagt cagcctg

17

<210> 112

<211> 23

<212> DNA

<213> HUMAN

<400> 112

aggcttgctc ctccccacc cag

23

<210> 113

<211> 18

<212> DNA

<213> HUMAN

<400> 113

agactttggc tccatctc

18

<210> 114

<211> 20

<212> DNA

<213> HUMAN

<400> 114

cactccaccc ccatacctagc

20

<210> 115

<211> 26

<212> DNA

<213> HUMAN

<400> 115

gggagagggc acagggccag acaaac

26

<210> 116

<211> 20

<212> PRT

<213> HUMAN

<400> 116

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly
1 5 10 15

Gly Gly Gly Ser
20

<210> 117

<211> 34

<212> DNA

<213> HUMAN

<400> 117

cgtctagaag gaattcagct agactggctc agca

34

<210> 118

<211> 15

<212> PRT

<213> HUMAN

<400> 118

Glu Val Lys Pro Lys Ser Glu Glu Glu Gly Ser Leu Lys Leu Glu

1

5

10

15

<210> 119

<211> 12

<212> PRT

<213> HUMAN

<400> 119

Gly Glu Glu Asp Leu Pro Gly Glu Glu Asp Leu Pro

1

5

10

<210> 120

<211> 12

<212> PRT

<213> HUMAN

<400> 120

Glu Glu Asp Leu Pro Gly Glu Glu Asp Leu Pro Gly

1

5

10

<210> 121

<211> 10

<212> PRT

<213> HUMAN

<400> 121

Glu Asp Leu Pro Gly Glu Glu Asp Leu Pro

1

5

10

<210> 122

<211> 12

<212> PRT

<213> HUMAN

<400> 122

Asp Leu Pro Gly Glu Glu Asp Leu Pro Gly Glu Glu
1 5 10

<210> 123

<211> 12

<212> PRT

<213> HUMAN

<400> 123

Leu Pro Gly Glu Glu Asp Leu Pro Gly Glu Glu Asp
1 5 10

<210> 124

<211> 12

<212> PRT

<213> HUMAN

<400> 124

Pro Gly Glu Glu Asp Leu Pro Gly Glu Glu Asp Leu
1 5 10

<210> 125

<211> 9

<212> PRT

<213> HUMAN

<400> 125

Ala Pro Gly Glu Glu Asp Leu Pro Ala
1 5

<210> 126

<211> 9

<212> PRT

<213> HUMAN

<400> 126

Ala Gly Glu Glu Asp Leu Pro Gly Ala
1 5

<210> 127

<211> 9

<212> PRT

<213> HUMAN

<400> 127

Ala Glu Glu Asp Leu Pro Gly Glu Ala

1

5

<210> 128

<211> 9

<212> PRT

<213> HUMAN

<400> 128

Ala Glu Asp Leu Pro Gly Glu Glu Ala

1

5

<210> 129

<211> 9

<212> PRT

<213> HUMAN

<400> 129

Ala Asp Leu Pro Gly Glu Glu Asp Ala

1

5

<210> 130

<211> 9

<212> PRT

<213> HUMAN

<400> 130

Ala Leu Pro Gly Glu Glu Asp Leu Ala

1

5

<210> 131

<211> 8

<212> PRT

<213> HUMAN

<400> 131

Ala Gly Glu Glu Asp Leu Pro Ala

1

5

<210> 132
<211> 8
<212> PRT
<213> HUMAN

<400> 132
Ala Glu Glu Asp Leu Pro Gly Ala
1 5

<210> 133
<211> 8
<212> PRT
<213> HUMAN

<400> 133
Ala Glu Asp Leu Pro Gly Glu Ala
1 5

<210> 134
<211> 8
<212> PRT
<213> HUMAN

<400> 134
Ala Asp Leu Pro Gly Glu Glu Ala
1 5

<210> 135
<211> 8
<212> PRT
<213> HUMAN

<400> 135
Ala Leu Pro Gly Glu Glu Asp Ala
1 5

<210> 136
<211> 8
<212> PRT
<213> HUMAN

<400> 136
Ala Pro Gly Glu Glu Asp Leu Ala

1 5

<210> 137

<211> 9

<212> PRT

<213> HUMAN

<400> 137

Ala Lys Lys Met Lys Arg Arg Lys Ala

1 5

<210> 138

<211> 9

<212> PRT

<213> HUMAN

<400> 138

Ala Ile Thr Phe Asn Ala Gln Tyr Ala

1 5

<210> 139

<211> 9

<212> PRT

<213> HUMAN

<400> 139

Ala Ser Ala Ser Ala Pro Val Ser Ala

1 5

<210> 140

<211> 9

<212> PRT

<213> HUMAN

<400> 140

Ala Gly Gln Thr Arg Ser Pro Leu Ala

1 5

<210> 141

<211> 6

<212> PRT

<213> HUMAN

<400> 141

Ser Glu Glu Asp Ser Pro

1

5

<210> 142

<211> 6

<212> PRT

<213> HUMAN

<400> 142

Arg Glu Glu Asp Pro Pro

1

5

<210> 143

<211> 12

<212> DNA

<213> HUMAN

<400> 143

agggcacagg gc

12